

Customer	: ESRIN	Document Ref	: SST_CCI-PSD-UKMO-201
Contract No	: 4000109848/13/I-NB	Issue Date	: 11 April 2014
WP No	: 10	Issue	: 1

Project : SST-CCI- Phase-II

Title : Product Specification Document (PSD)

Abstract : This document contains the Product Specification for the ESA SST_cci project.

Author(s) : *Simon Good.*
N.R.

Checked by : *Chris Merchant*

Simon Good and Nick Rayner
Met Office

Chris Merchant
University of Reading
Science Leader

Accepted by : *Craig Donlon*

Craig Donlon
ESA Technical Officer
ESTEC

Distribution : SST_cci team members
ESA (Craig Donlon)

**EUROPEAN SPACE AGENCY
CONTRACT REPORT**

The work described in this report was done under ESA contract.
Responsibility for the contents resides in the author or organisation
that prepared it.



AMENDMENT RECORD

This document shall be amended by releasing a new edition of the document in its entirety. The Amendment Record Sheet below records the history and issue status of this document.

AMENDMENT RECORD SHEET

ISSUE	DATE	REASON FOR CHANGE
Phase-I		
1	01 Apr 2011	First issue
2	11 Nov 2011	Amended following comments from ESA (Craig Donlon)
3	15 Mar 2013	Amended by S. Good following comments from project team
Phase-II		
1	11 Apr 2014	Correction of issues identified in Phase-I of the project and update to show Phase-II products (Simon Good)

TABLE OF CONTENTS

1. INTRODUCTION.....	5
1.1 Purpose and Scope.....	5
1.2 Document revision control.....	6
1.3 Applicable Documents.....	6
1.4 Referenced Documents.....	6
1.5 Definitions of Terms.....	8
2. PRODUCT OVERVIEW.....	10
2.1 Summary of products.....	10
2.2 Improvements over existing data.....	15
3. FILE FORMAT AND METADATA.....	16
3.1 Rationale.....	16
3.2 Filename convention.....	17
3.3 Global attributes.....	20
3.4 Variable attributes.....	28
3.5 Unique product identifiers.....	32
4. PRODUCT SPECIFICATION.....	34
4.1 Format of the product description tables.....	34
4.2 Storage type definitions.....	35
4.3 L2P data.....	35
4.4 L3U data.....	67
4.5 L4 data.....	75
5. FILE SIZE ESTIMATES.....	85
6. ISSUES FOR FUTURE RELEASES OF THIS DOCUMENT.....	86
APPENDIX A: CDL LISTINGS OF NETCDF HEADERS.....	87
A.1 L2P example.....	87
A.2 L3U example.....	92
A.3 L4 example.....	97
APPENDIX B: SPECIFICATION OF FILES TO STORE SSTs FROM INDIVIDUAL RETRIEVALS.....	101
B.1 L2P auxiliary files.....	101
B.2 L3 auxiliary files.....	109
B.2.1 Auxiliary file size estimates.....	110
B.2.2 CDL listings of NetCDF headers for the auxiliary files.....	110

This Page Is Intentionally Blank

1. INTRODUCTION

1.1 Purpose and Scope

1.1.1 Purpose

This document provides descriptions and specifications of products that shall be produced by the sea surface temperature (SST) climate change initiative project (SST_cci) Phase-II, which is part of the European Space Agency (ESA) Climate Change Initiative (CCI). The specifications were originally developed from user requirements defined in the SST_cci User Requirements Document [RD.171] and have subsequently been revised following experience gained within the project and from users making use of the data. SST_cci Phase -II plans are to produce SST products as follows:

- Gridded, 0.05° resolution, daily, day-night separated reference SSTs from Along Track Scanning Radiometer (ATSR) satellite data.
- 4-25 km resolution swath-based SSTs from Advanced Very High Resolution Radiometer (AVHRR) satellite data.
- Analyses of the above two data streams (gridded at 0.05° resolution, daily).

The specifications contained in this Product Specification Document (PSD) provide a complete description of the products, including the definition of file level metadata. Specifications are defined for level 2, 3 and 4 data files. These are all based on the Group for High Resolution SST (GHRSSST) Data Specification version 2.0 (GDS2.0; document revision 2.007) [AD.1], are specified to be compatible with the climate and forecasting (CF) conventions [RD.177] and include the metadata specified by the CCI Data Standards Working Group (DSWG; [RD.215]).

1.1.2 Scope

This document covers the following aspects of requirement SST-TR-12 from the CCI Phase-I statement of work, annex G [RD.165] / SST-TR-25 from the CCI Phase-II statement of work, annex G [RD.369]:

- File metadata format and structure (Sections 2 and 4).
- Discovery metadata contained in the files and their structure (Sections 2 and 4)
- Long-term document revision control procedures (Section 1.2).
- Any other requirements relevant to SST ECV product specification (for example file format and naming convention; Sections 2.2 and 3.2).

This document does not include in its scope the analysis of the error budget for translating input data to ECV products [RD.164]. This is reported in a separate document [RD.174].

Information is included to allow the reader to trace the product specifications defined in this document back to the user requirements, both directly by referring to the user requirements document, [RD.171], and through performance metrics obtained by analysis of the user requirements as detailed in a technical note [RD.189].

This document is to be subjected to review by the international scientific community (SST-TR-13) [RD.165]. The main focus of this review will be discussions with the GHRSSST community.

1.2 Document revision control

Document revision control is necessary to manage any changes to the product specifications and to retain a permanent record of when and where updates were made. The following principles will be followed in order to ensure that this occurs:

- Every release of the product specification document should be archived in a central project archive. The files should be protected from accidental change or deletion and backed up to a separate location.
- For every new release the version number should be incremented. The document naming convention established for the SST_cci should be used to ensure a unique file name for every version of the document.
- A summary of changes to the document and the person responsible for making the changes should be recorded in the relevant place at the beginning of the document.

Each new release must be approved by the project science leader or a person nominated by the project science leader.

1.3 Applicable Documents

The following is a list of documents that must be read in conjunction with this document. Where referenced in the text, these are identified as AD.n, where 'n' is the number in the list below:

- AD.1 GHRSSST Science Team, cited 2010: The Recommended GHRSSST Data Specification (GDS) Revision 2.0 Technical Specifications. [Available online at http://www.ghrsst.org/modules/documents/documents/GDS2.0_TechnicalSpecifications_v2.0.pdf.]

1.4 Referenced Documents

The following is a list of documents with a bearing on the content of this report. Where referenced in the text, these are identified as RD.n, where 'n' is the number in the list below:

- RD.164 ESA Climate Change Initiative phase 1 – scientific user consultation and detailed specification – statement of work, Issue 1.4, Revision 1, 09/11/2009, Reference EOP-SEP/SOW/0031-09/SP; http://earth.eo.esa.int/workshops/esa_cci/ao6207SoW.pdf
- RD.165 Annex G to RD.164, Sea Surface Temperature ECV (SST_cci)
- RD.166 Taylor, K.E. and C. Doutriaux (2010), CMIP5 Model Output Requirements: File Contents and Format, Data Structure and Metadata; http://cmip-pcmdi.llnl.gov/cmip5/docs/CMIP5_output_metadata_requirements.pdf
- RD.167 Rew, R., G. Davis, S. Emmerson, H. Davies, E. Hartnett and D. Heimbigner (2010), The NetCDF Users Guide;

<http://www.unidata.ucar.edu/software/netcdf/docs/netcdf.pdf>

- RD.168 Climate Modelling User Group (2010), Requirement Baseline Document;
http://dialspace.dial.pipex.com/prod/dialspace/town/estate/gtp89/cmug/D1.2_U RD_v1.3.pdf
- RD.169 ESA Climate Change Initiative – CCI project guidelines, Issue 1, Revision 0,
05/11/2010, Reference EOP-DTEX-EOPS-SW-10-0002
- RD.170 NetCDF Attribute Convention on Dataset Discovery (ACDD):
<http://www.unidata.ucar.edu/software/netcdf-java/formats/DataDiscoveryAttConvention.html>
- RD.171 CCI Phase 1 (SST) (2010), User Requirements Document, Reference
SST_CCI-URD-UKMO-001; http://www.esa-sst-cci.org/sites/default/files/documents/admin/public/SST_cci%20URD%20UKMO-001%20Issue%202.pdf
- RD.172 CCI Phase 1 (SST), Data Access Requirements Document (in preparation)
- RD.173 CCI Phase 1 (SST), Product Validation Plan (in preparation)
- RD.174 CCI Phase 1 (SST), Error Characterisation (in preparation)
- RD.176 Olsen, L.M., G. Major, K. Shein, J. Scialdone, R. Vogel, S. Leicester, H. Weir, S. Ritz, T. Stevens, M. Meaux, C. Solomon, R. Bilodeau, M. Holland, T. Northcutt and R. A. Restrepo (2007), NASA/Global Change Master Directory (GCMD) Earth Science Keywords Version 6.0.0.0.0;
<http://gcmd.nasa.gov/Resources/valids/>
- RD.177 NetCDF Climate and Forecast (CF) Metadata Convention; <http://cf-pcmdi.llnl.gov/>
- RD.178 The UDUNITS-2 package;
<http://www.unidata.ucar.edu/software/udunits/udunits-2/udunits2.html>
- RD.189 CCI Phase 1 (SST): Technical note on product metrics derived from the user requirements document
- RD.215 Bennett, V. (2010), Format and Metadata Guidelines for CCI Data Products (draft).
- RD.219 Donlon, C.J. et al. (2009), The GODAE High Resolution Sea Surface Temperature Pilot Project, *Oceanography*, 22(3), 34-45.
- RD.369 ESA Climate Change Initiative Phase 2 Statement of Work, prepared by ESA Climate Office, Ref. CCI-PRGM-EOPS-SW-12-0012, Issue 1, Revision 2, June 7th, 2013

1.5 Definitions of Terms

The following terms have been used in this report with the meanings shown.

Term	Definition
AATSR	Advanced ATSR.
ACDD	Attribute Convention for Dataset Discovery.
ATSR	Along Track Scanning Radiometer; the ATSR series of instruments comprises ATSR-1, ATSR-2 and AATSR.
AVHRR	Advanced Very High Resolution Radiometer.
BEAM	Earth observation toolbox and development platform.
CDL	Network Common Data form Language.
CF	Climate and Forecasting [conventions for metadata].
CMIP	Climate Model Intercomparison Project.
DSWG	CCI Data Standards Working Group
ECMWF	European Centre for Medium-range Weather Forecasts.
ECV	Essential Climate Variable.
ERA-Interim	ECMWF reanalysis 'interim' product.
ERS	European Remote Sensing satellite series (ERS-1, ERS-2).
GAC	Global Area Coverage.
GCMD	Global Change Master Directory.
GDS	GHRSSST Data Specification.
GMPE	GHRSSST Multi-Product Ensemble
GHRSSST	Group for High Resolution SST.
HDF5	Hierarchical Data Format version 5
IDL	Interactive Data Language.
ISO	International Organisation for Standardisation
L1B	Level 1 B data; the data collected by the satellite sensor with processing (for example calibration) applied.
L2P	Level 2 pre-processed data; geophysical variables at the full resolution of the instrument, in a satellite projection with associated geographical information.
L3U	Level 3 uncollated data; gridded version of L2P data.

L3C	Level 3 collated data; L2P data from a single instrument that have been combined and mapped onto a space-time grid.
L3S	Level 3 super-collated data; L2P data from multiple instruments that have been combined and mapped onto a space-time grid.
L4	Level 4 data; a dataset formed by analysing lower level data to create a gridded product with no data gaps. The use of an infilling procedure to remove data gaps distinguishes L4 data from lower level data.
MATLAB	Matrix Laboratory
NetCDF	Network Common Data Format.
NWP	Numerical Weather Prediction.
Obs4MIPs	Observations for Model Intercomparison Projects.
OSI SAF	Ocean and Sea Ice Satellite Applications Facility.
OSTIA	Operational Sea surface Temperature and sea Ice Analysis.
RDAC	Regional Data Assembly Centre.
SSES	Single Sensor Error Statistics.
SST	Sea Surface Temperature.
SST_cci	SST climate change initiative [project].
SEVIRI	Spinning Enhanced Visible and Infrared Imager.
SLSTR	Sea and Land Surface Temperature Radiometer.
TBC	To be confirmed.
URL	Uniform Resource Locator.
UUID	Universally Unique Identifier.
WGS	World Geodetic System.
XML	Extensible Mark-up Language.

2. PRODUCT OVERVIEW

2.1 Summary of products

An overview of the products to be produced by the SST_cci Phase-II project is shown in Table 1. The table also records which satellite sensors will be used to create the products and the format of the input data. The final column indicates the level of data to be produced from each sensor and the resolution/grid spacing of the data in the files. The SST_cci Phase-II project is subdivided into three contractual phases (CPs). The time coverage of the products produced in each CP is indicated in Figure 1. It is planned that data produced in CP1 and CP3 will be made publicly available.

Table 1. Summary of SST_cci products, the source and level of data to be produced. L2 and L3 products contain data from a single sensor; L4 products will include data from both sensors.

Category of product and description	Satellite sensors & data to be used	Level of data to be produced for each sensor (resolution/grid spacing)
ATSR SSTs retrieved from data obtained from the ATSR series of sensors (comprising ATSR-1, ATSR-2, AATSR and, possibly within the lifetime of the project, SLSTR).	ATSR series (ATSR-1, ATSR-2, AATSR, SLSTR) (Envisat format)	L3U (0.05°)
AVHRR SSTs retrieved from data obtained from the AVHRR series of sensors.	AVHRR series global area coverage (GAC) data	L2P (4-25 km)
Analysis Gap-free daily analyses of the ATSR and AVHRR products.	The ATSR and AVHRR products described above.	L4 (0.05°)




Figure 1. Time spans of the products to be produced in each contractual phase (CP) of the SST_cci phase II project. The grey shading indicates the available time span of the ATSR and AVHRR satellite series. Dotted lines are used where the time span is yet to be finalised. SLSTR data from Sentinel-3 (not shown) may also become available.

The table indicates that the project will produce data at levels L2P, L3U and L4. These are all defined below. User requirements indicate that all will be useful to users (SST_CCI-UR-QUF-33, [RD.171]). As the project will receive some data at data level L1B, this is also defined below.

- L1B Level 1 B data; the data collected by the satellite sensor with processing (for example calibration) applied.
- L2P Level 2 pre-processed data; geophysical variables at the full resolution of the instrument, in a satellite projection with associated geographical information.
- L3U Level 3 uncollated data; gridded version of L2P data.
- L4 Level 4 data; a dataset formed by analysing lower level data to create a gridded product with no data gaps. The use of an infilling procedure to remove data gaps distinguishes L4 data from lower level data.

Each entry in Table 1 is expanded in Table 2. This table provides full details of each product. This includes aspects such as the file names, the file sizes, the target for stability etc.

The headings to the columns in Table 2 define unique names for each product. These shall be appended in brackets with the name of an individual sensor if it is necessary to distinguish between sensors in a series. For example the individual components of the ATSR product shall be referred to as ATSR (ATSR-1), ATSR (ATSR-2) and ATSR (AATSR).

Table 2. Detailed description of the SST_cci products. The column headings define unique names for each product. Where it is necessary to distinguish between data from individual sensors in products that contain a series of sensors the name of the sensor shall be appended to the product name in brackets. For example the ATSR-1 part of the ATSR product shall be referred to as ATSR (ATSR-1). File size estimates are based on the following grid sizes: AVHRR L2P – 625 × 7000, L3 and L4 data - 7,200 × 3,600 and one time step per file. User requirements for each aspect of the products are included and are referenced as codes beginning with SST_CCI-UR-; full details of these requirements are in [RD.171].

Product	ATSR	AVHRR	Analysis
Satellite sensors used	ATSR series (ATSR-1, ATSR-2, AATSR, SLSTR if available)	AVHRR series GAC data	ATSR series and AVHRR series
Satellite sensor type	Infrared	Infrared	Infrared
Satellite platform	ERS-1, ERS-2, Envisat, Sentinel-3	NOAA platforms, Metop-A	ERS-1, ERS-2, Envisat, Sentinel-3, NOAA platforms, Metop-A
Satellite orbit	Polar	Polar	Polar
Input data source	ESA	NOAA	SST_cci (ATSR and AVHRR products)
Input data format	Envisat format	AVHRR GAC	NetCDF compliant with this document
Input data level	L1B	L1B	L3U (ATSR) and L2P (AVHRR)
Input data frequency	One file per orbit per sensor (typically 14 orbits per day, one sensor operating)	One file per orbit per sensor (typically 14 orbits per day, more than one sensor operating)	Typically 28 orbits per day, one file per orbit
Input data grid	0.05° grid	4-25 km spaced irregular grid	0.05° grid (ATSR) and 4-25 km irregular grid (AVHRR)
Input data spatial coverage	Global	Global	Global
Processing to be applied by the project	SST retrieval; cloud and ice screening; depth and time adjustment to SSTs; uncertainty estimates	SST retrieval; cloud and ice screening; depth and time adjustment to SSTs; uncertainty estimates	Analysis

Product	ATSR	AVHRR	Analysis
Output data level	L3U	L2P	L4
Output data level user requirement	SST_CCI-UR-QUF-33: indicates that all the levels of data to be produced by the project (levels 2, 3 and 4) will be useful to users.		
Output data frequency	One file per orbit per sensor (typically 14 orbits per day, one sensor operating)	One file per orbit per sensor (typically 14 orbits per day, more than one sensor operating)	One file per day
Output data frequency user requirement	SST_CCI-UR-QUF-38: the most common requirements for data frequency at a location are monthly (threshold), daily (breakthrough) and 3 hourly (objective). See also [RD.189] Tables 4 and 19.		
Output grid	0.05° grid	4-25 km spaced irregular grid	0.05° grid
Output grid user requirement	SST_CCI-UR-QUF-36: 0.05° exceeds the user requirement at breakthrough level (i.e. data at this grid spacing will result in a significant improvement to users' applications) and is sufficient to allow the data to be used for a high proportion of applications [RD.189]. See also [RD.189] Tables 3 and 18.		
File size estimate (assuming no compression)	674 MB	140 MB	181 MB
File size user requirement	SST_CCI-UR-QUF-87: requirement for maximum file size varies considerably between users, between <100 KB and >10 GB. See also [RD.189] Table 33.		
Output file format	CF compliant NetCDF		
Output file format user requirement	SST_CCI-UR-QUF-89: CF compliant NetCDF is required.		
Spatial coverage	Global		
Spatial coverage user requirement	SST_CCI-UR-QUF-42: global coverage is required.		
Product start date	CP1: 08/1991 CP2: 08/1991 (TBC) CP3: 08/1991	CP1: 09/1991 CP2: 1981 CP3: 1981	CP1: None CP2: TBC CP3: 1981
Product end date	CP1: 04/2012 CP2: 1996 (TBC) CP3: 04/2012	CP1: 2012-13 (TBC) CP2: 1996 CP3: 2016	CP1: None CP2: TBC CP3: 2016

Product	ATSR	AVHRR	Analysis
Product length	CP1: approx. 21 years CP2: approx. 5 years CP3: approx. 21 years	CP1: 21-22 years CP2: 16 years CP3: 36 years	CP1: 0 CP2: 4 years CP3: 36 years
Product length user requirement	SST_CCI-UR-QUF-43: there is a clear requirement for data records longer than 30 years. See also [RD.189] Tables 8 and 23.		
SST depth	Skin and 20 cm	Skin and 20 cm	20 cm
SST depth user requirement	SST_CCI-UR-QUF-46: skin SST is the most common requirement, followed by SSTs at depths roughly corresponding to the range of traditional in situ observations (20 cm and 5 m). See also [RD.189] Tables 17 and 32.		
SST time of day	Platform dependent time and adjusted to 10.30 am and pm local time	Platform dependent time and adjusted to 10.30 am and pm local time	Representative of daily average
SST time of day user requirement	SST_CCI-UR-QUF-41: SSTs are most commonly required at synoptic observing times. See also [RD.189] Tables 5 and 20. However this does not match well with the observing system. SST_CCI-UR-QUF-40: Temporal averages are acceptable to the majority of users – the adjusted SSTs and analyses will be representative of a daily average.		
Target accuracy	0.1°C when evaluated over length scales of ≤ 1000 km		
Acceptable level of bias user requirement	SST_CCI-UR-QUF-48: the highest acceptable bias level is 0.1°C, and, SST_CCI-UR-QUF-49: 0.1°C is the required precision. While the user requirement is that the achievement of both should be demonstrated over spatial scales of 100 km, the achievable scales are controlled by the density of comparison data, and will usually be longer than this. See also [RD.189] Tables 9, 10, 11, 12, 24, 25, 26, 27.		
Target stability	0.1°C/decade		
Stability user requirement	SST_CCI-UR-QUF-50: 0.1°C per decade, to be demonstrated on spatial scales of 100 km. See also [RD.189] Tables 13, 14, 28, 29.		
Ancillary data to be included	Surface wind speed from the ERA-Interim reanalysis	Sea ice concentration and uncertainties; taken from OSI SAF data	
Ancillary data user requirement	SST_CCI-UR-QUF-70		SST_CCI-UR-QUF-69
File names – read from top to bottom to obtain the file name	<Indicative Date><Indicative Time>-ESACCI-		
	L3U	L2P	L4

Product	ATSR	AVHRR	Analysis
for each product	_GHRSSST-		
Definitions of the fields contained in chevrons (<>) are in Section 3.2.	SSTskin-[ATSR1, ATSR2 or AATSR]	SSTskin-AVHRR<X>_G	SSTdepth-OSTIA-GLOB
	-v2.0-fv<FileVersion>.nc		

2.2 Improvements over existing data

The new products to be produced by the SST_cci project will address a number of the weaknesses of existing SST data that are perceived by users and take advantage of their strengths. These strengths and weaknesses are described in full in [RD.171]; some points of relevance to the SST_cci products are summarised below:

- Long term records of SST are available from the ATSR and AVHRR series of sensors. ATSR series data are viewed as weak in spatial sampling and the frequency of observations. However, its bias characteristics and stability are seen as strengths. Frequency of observations is seen as a strength of AVHRR series data but its bias characteristics received mixed responses from users.
 - By using new algorithms and cross referencing the AVHRR data to the ATSR data the SST_cci project produces SST products that have the bias characteristics and stability approaching that of ATSR alone with the coverage of AVHRR.
- A >30 year climate data record is required for many applications.
 - The new SST_cci phase II products will extend those produced in phase I of the project to provide a >30 year climate data record.
- Characterisation of uncertainties is generally viewed as being weak.
 - The project improves this situation by providing uncertainty estimates split into components with different spatial and temporal correlation characteristics.

3. FILE FORMAT AND METADATA

3.1 Rationale

The format of the data files shall be Network Common Data Format (NetCDF) and shall follow Climate and Forecasting (CF) conventions. This is the format preferred by the majority of potential users of the products (user requirement SST_CCI-UR-QUF-89, [RD.171]). The use of NetCDF also allows the use of a variety of different tools/languages to manipulate and view the data, for example BEAM and the tools available within languages such as the Interactive Data Language (IDL), the Matrix Laboratory (MATLAB) and Python.

CF compliant NetCDF files are already very well established in the SST user community owing to the success of the GHRSSST project ([RD.219]; <https://www.ghrsst.org/>). GHRSSST provides SST data products in a CF compliant NetCDF data format, the latest version being the GDS2.0 [AD.1]. The GDS2.0 already includes much of the data and metadata that users require and that SST_cci wish to include in the files, but also has the flexibility of allowing additional data fields to be added as required. The user requirements gathered by the SST_cci project shows some limited support for the GDS2.0 for the NetCDF files (SST_CCI-UR-QUF-89, [RD.171]). It is not clear from the results whether this limited support is due to lack of awareness of the specification or a desire for an alternative. Adopting the GDS2.0 specification would provide benefits to users as some will already have experience with the files and may have existing software and tools to work with the data. It will also ensure easy incorporation of the SST_cci products into the GHRSSST framework. ESA have mandated use of GDS2.0 for this project.

It would also be desirable for the SST_cci data to be available in a form that is familiar to climate modellers to allow ease of comparison of observational data with climate model data [RD.168]. Format specifications exist to allow this ('Obs4MIPS format'; see <http://obs4mips.llnl.gov/>), However, this has some differences to the way that the storage of data in NetCDF format is implemented by GHRSSST. These differences mean that it is not possible to produce data files that are consistent with both specifications. The SST_cci project shall produce files that are consistent with the GDS2.0 specification as described in this document. Files in the Obs4MIPS format will be produced from the data output in CP3.

The format of NetCDF used for GDS2.0 data files is preferentially 'NetCDF-4 classic' and therefore this shall be used for SST_cci files. The term 'NetCDF-4 classic' combines two aspects of the files: the underlying file format and the data model. These are explained briefly below:

Three different underlying formats could be used:

- 'classic': this is compatible with all versions of the NetCDF library;
- '64-bit offset': this allows larger datasets to be written but is not compatible with versions of the NetCDF library prior to 3.6.0;
- NetCDF-4; this uses Hierarchical Data Format version 5 (HDF5) as its base format and gives access to desirable features such as internal per variable compression; it is compatible with the NetCDF library from version 4.

Although the NetCDF-4/HDF5 format is not compatible with older versions of the library, the additional features such as its compression abilities gives clear advantages over the other options.

There are two NetCDF data models: 'classic' and 'enhanced' [RD.167]. The classic data model has a number of limitations compared to enhanced, for example the enhanced data model includes a greater number of allowed storage types. Either can be written using the NetCDF-4/HDF5 underlying format. The advantage of using the classic model is that it maintains compatibility with legacy software.

In summary:

- The SST_cci files shall be internally compressed NetCDF-4/HDF5 format and shall use the NetCDF classic data model.
- The file specifications shall follow CF conventions.
- The files will be based on and be consistent with the GHRSSST GDS2.0 [AD.1].

3.2 Filename convention

SST_cci shall follow the file naming convention specified in [AD.1], i.e.:

<Indicative Date><Indicative Time>-<RDAC>-<Processing Level>_GHRSSST-<SST Type>-<Product String>-<Additional Segregator>-v<GDS Version>-fv<FileVersion>.<File Type>

The components to this list are summarised in [AD.1] Table 7-1 with further explanatory information in the text of that document. That table is reproduced here in Table 3 with an additional column (headed SST_cci definition) that details the implementation of the filenaming convention that shall be used by the SST_cci project. All the information required to define the SST_cci files is contained within this document. However, note that references in the Description column of the table are to sections in the [AD.1] document and these shall be referred to if further explanation is desired. In places, clarifications to the [AD.1] version of the text have been made and these are indicated in italics.

Table 3. Components of the filenames; based on [AD.1] section 7.1 – refer there for more details. The final column details the implementation for the SST_cci project. In the Description column the references refer to sections of [AD.1]. All references in the SST_cci column are to parts of this document.

Name	Definition	Description	SST_cci definition
<Indicative Date>	YYYYMMDD	The identifying date for this data set. See Section 7.2 in [AD.1]. The section states that: YYYY is the four-digit year, MM is the two-digit month from 01 to 12, and DD is the two-digit day of month from 01 to 31 and The date used should best represent the observation date for the dataset.	As stated in Description column.
<Indicative Time>	HHMMSS	The identifying time for this data set. The time used is dependent on the <Processing Level> of the data set: L2P: start time of granule L3U: start time of granule L3C and L3S: centre time of the collation window L4 and GMPE: nominal time of analysis All times should be given in UTC. See Section 7.3 in [AD.1]. The section states that: HH is the two-digit hour from 00 to 23, MM is the two-digit minute from 00 to 59, and SS is the two-digit second from 00 to 59 and All times should be given in UTC and should be chosen to best represent the observation time for this dataset. Note: RDACs should ensure the applications they use to determine UTC properly account for leap seconds.	As stated in Description column.
<RDAC>	The RDAC where the file was created.	The Regional Data Assembly Centre (RDAC) code.	For the SST_cci products, this is: ESACCI
<Processing Level>	The data processing level code (L2P, L3U, L3C, L3S, or L4)	The data processing level code, defined in Section 7.5 in [AD.1].	See Table 1 in this document.
<SST type>	The type of SST data included in the file.	Conforms to the GHRSSST definitions for SST, defined in Section 7.6 in [AD.1].	This should be set to be the primary SST stored in the file. See Table 4.

Name	Definition	Description	SST_cci definition
<Product String>	A character string identifying the SST product set. The string is used uniquely within an RDAC but may be shared across RDACs.	The unique “name” within an RDAC of the product line. See Section 7.7 <i>in [AD.1]</i> for the product string lists, one each for L2P, L3, L4, and GMPE products. See Section 7.7 <i>in [AD.1]</i> .	See Table 5 in this document.
<Additional Segregator>	Optional text to distinguish between files with the same <Product String>. Dashes are not allowed within this element.	This text is used since the other filename components are sometimes insufficient to uniquely identify a file. For example, in L2P or L3U (un-collated) products this is often the original file name or processing algorithm. Note, underscores should be used, not dashes. For L4 files, this element should begin with the appropriate regional code as defined in Section 7.8. This component is optional but must be used in those cases where non-unique filenames would otherwise result.	For L2 and L3 products this is not used. For L4 products it is GLOB (GLOB is required to indicate spatial coverage)
<GDS Version>	nn.n	Version number of the GDS used to process the file. For example, GDS 2.0 = “02.0”.	To be set to “02.0”.
<File Version>	xx.x	Version number for the file, for example, “01.0”.	As stated in Description column.
<File Type>	NetCDF data file suffix (nc) or ISO metadata file suffix (xml).	Indicates this is a NetCDF file containing data or its corresponding ISO-19115 metadata record in XML.	Will be nc for SST_cci files.

Table 4. Values and definitions of <SST type>. Based on [AD.1] section 7.6 – refer there for more details.

SST type	Description
SSTskin	SST sensed by infrared instruments (at depth of approximately 10 µm).
SSTsubskin	SST sensed by microwave instruments (at depth of approximately 1 mm).
SSTdepth	Temperature at a specific depth. The depth shall be included in the file metadata.
SSTfnd	Temperature from which the diurnal thermocline develops each day.

Table 5: Product strings for data to be processed; based on [AD.1] Tables 7-5 to 7-7 – refer there for more details. Codes for SLSTR should be confirmed with the GHRSSST project before use.

Sensor/analysis	Platform (if applicable)	Product String	Comment
ATSR	ERS-1	ATSR1	
ATSR	ERS-2	ATSR2	
AATSR	Envisat	AATSR	
AVHRR_GAC	NOAA-<X> or MetOpA	AVHRR<X>_G	<X> is <i>either</i> : if the satellite hosting the AVHRR is a NOAA platform, <X> is the satellite number, <i>or</i> , if the AVHRR is on Metop-A, <X> is 'MTA'.
OSTIA		OSTIA	L4 analysis using the OSTIA system
SLSTR	SENTINEL_3A	S3A_SLSTR	TBC

3.3 Global attributes

Attributes are metadata that are stored in a NetCDF file. Global attributes contain information that applies to the whole contents of the NetCDF file.

All GHRSSST format data files share the same set of global attributes. These are listed in [AD.1] Table 8-1. The CCI DSWG also specify a set of global attributes that should be included in the SST_cci data files([RD.215]), some of which are not in the GHRSSST specification. In addition an attribute called “creator_processing_institution” has been included to indicate the institution that created products as part of the SST_cci project

Table 6 contains a list of the global attributes. Much of this is reproduced from Table 8-1 in [AD.1]. However, annotations to the table have been made in places in order to clarify the text; these are indicated by italics. The additional attributes specified by the CCI DSWG are also included. These are the entries that include ‘CCI DSWG’ in the Source column of the table. The “creator_processing_institute” attribute has SST_cci within the Source column. Other points to note are:

- The definition of the shading of the rows in the table is given in Table 7.
- All references in the Description column refer to the contents of [AD.1]. Although all the information required to create the SST_cci files is contained here, these can be used to obtain more information about GDS2.0 files.
- The final column, headed ‘SST_cci definition’ contains information about the implementation of the attributes for the SST_cci files. All references in this column are to parts of this document.

As revealed by the column headed ‘Source’, the definition of the attributes was based on CF conventions, the Unidata Attribute Convention for Dataset Discovery (ACDD)

[RD.170], the requirements of the GHRSSST project [AD.1] and the specifications provided by the CCI DSWG.

Table 6. Global attributes that are included in all GHRSSST format data files and additional attributes specified by the CCI DSWG. Much of the contents of this table is reproduced from [AD.1] and further information can be found there. Annotations and changes to the original text are made in places and these are written in italics. The meaning of the colours is given in Table 7. Note that attributes described as having the format ‘string’ shall actually be stored in the NetCDF file as an array of characters. References in the Description column are to parts of [AD.1]; references in the SST_cci definition column are to parts of this document.

Global Attribute Name	Format	Description	Source	SST_cci definition
Conventions	string	A text string identifying the netCDF conventions followed. This attribute should be set to the version of CF used and should also include the ACDD. For example: "CF-1.4, Unidata Observation Dataset v1.0".	CF	CF-1.5, Unidata Observation Dataset v1.0
title	string	A descriptive title for the GHRSSST data set	CF, ACDD	See examples in Section 4 and Appendix A.
summary	string	A paragraph describing the dataset.	ACDD	See examples in Section 4 and Appendix A.
references	string	Published or web-based references that describe the data or methods used to produce it.	CF	Include any relevant publications or webpages.
institution	string	GHRSSST RDAC code where the data were produced. See Table 7-2 in [AD.1] for available codes (<i>this version of the GDS does not, however, include the SST_cci RDAC code</i>).	CF, ACDD	The RDAC code for the SST_cci products is ESACCI
history	string	History of all applications that have modified the original data to create this file.	CF, ACDD	Appendix A contains a basic example of this.

Global Attribute Name	Format	Description	Source	SST_cci definition
comment	string	Miscellaneous information about the data or methods used to produce it.	CF, ACDD	Used to indicate the centre where the data were produced. Set to: These data were produced at <institution> as part of the ESA SST CCI project.
license	string	Describe any restrictions to data access, use, and distribution. GHR SST data sets should be freely and openly available to comply with the R/GTS framework, with no restrictions. However, if a user should submit a simple registration via a web form, for example, the URL could be given here. Default to "GHR SST protocol describes data use as free and open."	ACDD	Creative Commons Licence by attribution (http://creativecommons.org/licenses/by/3.0/)
id	string	The unique GHR SST character string for this product. All GHR SST SST products have one, and they are listed in Table 7-10 in [AD.1].	ACDD	See Section 3.5 for the list of strings to be used by the SST_cci project.
naming_authority	string	Fixed as "org.ghrsst" following ACDD convention.	ACDD	org.ghrsst
product_version	string	The product version of this data file, which may be different than the file version used in the file naming convention (Section 7 in [AD.1]).	GDS	See Section 3.2 of this document
uuid	string	A Universally Unique Identifier (UUID). Numerous, simple tools can be used to create a UUID, which is inserted as the value of this attribute. See http://en.wikipedia.org/wiki/Universally_Unique_Identifier for more information and tools.	GDS	[RD.169] suggests the use of http://www.oss.org/pkg/lib/uuid/ , and this will be followed.

Global Attribute Name	Format	Description	Source	SST_cci definition
tracking_id	string	Identical to uuid (i.e. this shall contain the same value as uuid).	CCI DSWG	As stated in Description column. This is an alternative name for the UUID as used in the CMIP5 specification [RD.166].
gds_version_id	string	GDS version used to create this data file. For example, "2.0".	GDS	The GDS version used for the SST_cci product specification was 2.0.
netcdf_version_id	string	Version of netCDF libraries used to create this file. For example, "4.1.1"	GDS	As stated in Description column. Optionally can also include the date for example "4.2.1.1 of Oct 19 2012 14:25:16"
date_created	string	The date and time the data file was created in the form "yyyymmddThhmmssZ". This time format is ISO 8601 compliant.	ACDD	As stated in Description column.
file_quality_level	integer	A code value: 0 = unknown quality 1 = extremely suspect (frequent problems, e.g. with known satellite problems) 2 = suspect (occasional problems, e.g. after launch) 3 = excellent (no known problems)	GDS	As stated in Description column.

Global Attribute Name	Format	Description	Source	SST_cci definition
spatial_resolution	string	A string describing the approximate resolution of the product. For example, "1.1km at nadir"	GDS	Value depends on the product. See Table 1 or Table 2.
start_time	string	Date and time of the first measurement in the data file in the form "yyyymmddThhmmssZ". This time format is ISO 8601 compliant.	GDS	As stated in Description column.
time_coverage_start	string	Identical to start_time. Included for increased ACDD compliance.	ACDD	As stated in Description column.
stop_time	string	Date and time of the last measurement in the data file in the form "yyyymmddThhmmssZ". This time format is ISO 8601 compliant.	GDS	As stated in Description column.
time_coverage_end	string	Identical to stop_time. Included for increased ACDD compliance.	ACDD	As stated in Description column.
time_coverage_duration	string	Duration of time coverage in ISO 8601 compliant format; the difference between time_coverage_end and time_coverage_start.	CCI DSWG, ACDD	In the form PdDThHmMsS or other ISO 8601 compliant format where d is the number of days, h is the number of hours, m is the number of minutes, s is the number of seconds, omitting dD etc. if the number is zero.
time_coverage_resolution	string	Time resolution of the data in the file in ISO 8601 compliant format.	CCI DSWG, ACDD	Orbit repeat period (for single orbit files) or frequency of data files.

Global Attribute Name	Format	Description	Source	SST_cci definition
northernmost_latitude	float	Decimal degrees north, range - 90 to +90. This is equivalent to ACDD geospatial_lat_max.	GDS	As stated in Description column; see also examples in Appendix A.
geospatial_lat_max	float	Identical to northernmost_latitude.	CCI DSWG, ACDD	As stated in Description column.
southernmost_latitude	float	Decimal degrees north, range - 90 to +90. This is equivalent to ACDD geospatial_lat_min.	GDS	As stated in Description column; see also examples in Appendix A.
geospatial_lat_min	float	Identical to southernmost_latitude.	CCI DSWG, ACDD	As stated in Description column.
easternmost_longitude	float	Decimal degrees east, range - 180 to +180. This is equivalent to ACDD geospatial_lon_max.	GDS	As stated in Description column; see also examples in Appendix A.
geospatial_lon_max	float	Identical to easternmost_longitude.	CCI DSWG, ACDD	As stated in Description column.
westernmost_longitude	float	Decimal degrees east, range - 180 to +180. This is equivalent to ACDD geospatial_lon_min.	GDS	As stated in Description column; see also examples in Appendix A.
geospatial_lon_min	float	Identical to westernmost_longitude.	CCI DSWG, ACDD	As stated in Description column.
geospatial_vertical_min	float	Minimum of the vertical extent of the data; positive values indicate distances above the surface.	CCI DSWG, ACDD	To be set to the representative depth of the deepest SST in the file.
geospatial_vertical_max	float	Maximum of the vertical extent of the data; positive values indicate distances above the surface.	CCI DSWG, ACDD	To be set to the representative depth of the shallowest SST in the file.

Global Attribute Name	Format	Description	Source	SST_cci definition
source	string	Comma separated list of all source data present in this file. List SST sources first, followed by Auxiliary sources. If the source is a GHR SST product, use the GHR SST unique string listed in Table 7-10. For other sources, following the best practice described in Section 7.9 in [AD.1].	CF	See Section 3.5 in this document for the source codes relevant to the SST_cci products.
platform	string	Satellite(s) used to create this data file. Select from the entries found in the Satellite Platform column of Table 7-5 in [AD.1] and provide as a comma separated list if there is more than one.	GDS	See Table 5 in this document for the platforms relevant to the SST_cci products.
sensor	string	Sensor(s) used to create this data file. Select from the entries found in the Satellite Sensor column of Table 7-5 in [AD.1] and provide as a comma separated list if there is more than one.	GDS	See Table 5 in this document for the platforms relevant to the SST_cci products.
Metadata_Conventions	string	Unidata Dataset Discovery v1.0	ACDD	As stated in Description column.
metadata_link	string	Link to collection metadata record at archive	ACDD	http://www.esa-cci.org
keywords	string	Typically GCMD Science Keyword: "Oceans > Ocean Temperature > Sea Surface Temperature"	ACDD	As stated in Description column.
keywords_vocabulary	string	"NASA Global Change Master Directory (GCMD) Science Keywords" as defined in [RD.176].	ACDD	As stated in Description column.
standard_name_vocabulary	string	"NetCDF Climate and Forecast (CF) Metadata Convention"	ACDD	As stated in Description column.
geospatial_lat_units	string	Units of the latitudinal resolution. Typically "degrees_north"	ACDD	degrees_north

Global Attribute Name	Format	Description	Source	SST_cci definition
geospatial_lat_resolution	float	Latitude Resolution in units matching geospatial_lat_units.	ACDD	See examples in Appendix A.
geospatial_lon_units	string	Units of the longitudinal resolution. Typically "degrees_east"	ACDD	degrees_east
geospatial_lon_resolution	float	Longitude Resolution in units matching geospatial_lon_resolution.	ACDD	See examples in Appendix A.
acknowledgment	string	Information about funding source and how to cite the use of these data.	ACDD	Funded by ESA.
creator_name	string	Provide a name and email address for the most relevant point of contact at the producing RDAC, as well as a URL relevant to this data set.	ACDD	SST_cci
creator_email	string		ACDD	science.leader@esa-sst-cci.org
creator_url	string		ACDD	http://www.esa-sst-cci.org
creator_processing_institution	string	Contains additional information about the institution that created the product within the CCI project.	SST_cci	These data were produced at <institution> as part of the ESA SST CCI project.
project	string	"Group for High Resolution Sea Surface Temperature"	ACDD	Climate Change Initiative – European Space Agency
publisher_name	string	<i>The name of the data publisher; for GHRSSST data this is The GHRSSST Project Office</i>	ACDD	ESACCI
publisher_url	string	<i>The web address of the data published; for GRHSST data this is http://www.ghrsst.org</i>	ACDD	http://www.esa-sst-cci.org
publisher_email	string	<i>The email address of the data publisher; for GHRSSST data this is ghrsst-po@nceo.ac.uk</i>	ACDD	science.leader@esa-SST-cci.org

Global Attribute Name	Format	Description	Source	SST_cci definition
processing_level	string	GHRSSST definitions are the options: L2P, L3U, L3C, L3S, L4 and GMPE	ACDD	Value depends on the product. See Table 1 or Table 2.
cdm_data_type	string	“swath” or “grid”	ACDD, GDS	“swath” if L2P file, otherwise “grid”
product_specification_version	string	The file name (excluding extension) of the product specification document to which the file relates.	SST_cci	For example SST_CCI-PSD-UKMO-201-Issue-1-signed

Table 7. Key to colours used in Table 6 and Table 8. Based on Table 4-2 in [AD.1].

Colour	Meaning
Blue	Mandatory item.
Violet	An item that is mandatory for only certain situations.

3.4 Variable attributes

In the context of a NetCDF file a variable refers to an array of data stored within the file. Each variable can have attributes associated with it. [AD.1] defines attributes that should be provided with variables. As for the global attributes, the table of variable attributes from [AD.1] (Table 8-2) is reproduced here (Table 8). Annotations to the original text are indicated by italics. References within the table in the Description column refer to the contents of [AD.1] or to other external references. The SST_cci implementation of the variables is given in the column headed ‘SST_cci definition’. All references in this column are to parts of this document. There are some differences to the attributes to be used that depends on the nature of the data being stored; these are identified using violet shading (see Table 7).

Table 8. List of variable attributes. The contents of the first four columns of this table are reproduced from [AD.1] and further information can be found there. References in those columns refer to parts of [AD.1]. Annotations and changes to the [AD.1] text are identified by italics. The final column contains the SST_cci implementation of the attributes. All references in that column refer to parts of this document. The meaning of the colours is given in Table 7. Note that attributes described as having the format 'string' shall actually be stored in the NetCDF file as an array of characters.

Variable attribute name	Format	Description	Source	SST_cci definition
_FillValue	Must be same as the variable type	A value used to indicate array elements containing no valid data. This value must be of the same type as the storage (packed) type; should be set as the minimum value for this type. Note that some netCDF readers are unable to cope with signed bytes and may, in these cases, report fill as 128. Some cases will be reported as unsigned bytes 0 to 255.	CF	As stated in Description column.
units	string	Text description of the units, preferably S.I., and must be compatible with the Unidata UDUNITS-2 package [RD.178]. For a given variable (e.g. wind speed), these must be the same for each dataset.	CF, ACDD	See specifications in Section 4.
scale_factor	Must be expressed in the unpacked data type	To be multiplied by the variable to recover the original value. Defined by the producing RDAC. Valid values within valid_min and valid_max should be transformed by scale_factor and add_offset, otherwise skipped to avoid floating point errors.	CF	See specifications in Section 4. The type of this attribute defines the type of the unpacked data.
add_offset	Must be expressed in the unpacked data type	To be added to the variable after multiplying by the scale factor to recover the original value. If only one of scale_factor or add_offset is needed, then both should be included anyway to avoid ambiguity, with scale_factor defaulting to 1.0 and add_offset defaulting to 0.0. Defined by the producing RDAC.	CF	See specifications in Section 4. The type of this attribute defines the type of the unpacked data.
long_name	string	A free-text descriptive variable name.	CF, ACDD	See specifications in Section 4.

Variable attribute name	Format	Description	Source	SST_cci definition
valid_min	Expressed in same data type as variable	Minimum valid value for this variable once they are packed (in storage type). The fill value should be outside this valid range. Note that some netCDF readers are unable to cope with signed bytes and may, in these cases, report valid min as 129. Some cases as unsigned bytes 0 to 255. Values outside of valid_min and valid_max will be treated as missing values.	CF	See specifications in Section 4.
valid_max	Expressed in same data type as variable	Maximum valid value for this variable once they are packed (in storage type). The fill value should be outside this valid range. Note that some netCDF readers are unable to cope with signed bytes and may, in these cases, report valid min as 127.	CF	See specifications in Section 4.
standard_name	string	Where defined, a standard and unique description of a physical quantity. For the complete list of standard name strings, see [RD.177]. Do not include this attribute if no standard_name exists.	CF, ACDD	See specifications in Section 4.
comment	string	Miscellaneous information about the data or the methods used to produce it.	CF	As stated in Description column. See examples in Appendix A.
source	string	<p>For L2P and L3 files: For a data variable with a single source, use the GHRSSST unique string listed in Table 7-10 in [AD.1] if the source is a GHRSSST SST product. For other sources, following the best practice described in Section 7.9 in [AD.1] to create the character string.</p> <p>If the data variable contains multiple sources, set this string to be the relevant “sources of” variable name. For example, if multiple wind speed sources are used, set source = sources_of_wind_speed.</p> <p>For L4 and GMPE files: follow the source convention used for the global attribute of the same name, but provide in the comma-separated list</p>	CF	<p>String containing the source of data, or comma separated list if there are multiple sources.</p> <p>Section 3.5 details the sources of data to be used by the SST_cci products and the names for these that shall be used</p>

Variable attribute name	Format	Description	Source	SST_cci definition
		only the sources relevant to this variable.		to fill this attribute.
references	string	Published or web-based references that describe the data or methods used to produce it. Note that while at least one reference is required in the global attributes (See Table 8-1 in [RD.87], references to this specific data variable may also be given.	CF	As stated in Description column. Note that the equivalent in this document to the table that is referred to is Table 6.
axis	string	For use with coordinate variables only. The attribute 'axis' may be attached to a coordinate variable and given one of the values "X", "Y", "Z", or "T", which stand for a longitude, latitude, vertical, or time axis respectively. See: http://cf-pcmdi.llnl.gov/documents/cf-conventions/1.4/cf-conventions.html .	CF	See specifications in Section 4.
positive	string	For use with a vertical coordinate variables only. May have the value "up" or "down". For example, if an oceanographic netCDF file encodes the depth of the surface as 0 and the depth of 1000 meters as 1000 then the axis would set positive to "down". If a depth of 1000 meters was encoded as -1000, then positive would be set to "up". See the section on vertical-coordinate in <i>the CF conventions document</i> [RD.177].	CF	This is not used in SST_cci files.
coordinates	string	Identifies auxiliary coordinate variables, label variables, and alternative coordinate variables. See the section on coordinate-system in <i>the CF conventions document</i> [RD.177].	CF	As stated in Description column.

Variable attribute name	Format	Description	Source	SST_cci definition
grid_mapping	string	Use this for data variables that are on a projected grid. The attribute takes a string value that is the name of another variable in the file that provides the description of the mapping via a collection of attached attributes. That named variable is called a grid mapping variable and is of arbitrary type since it contains no data. Its purpose is to act as a container for the attributes that define the mapping. See the section on mappings-and-projections in <i>the CF conventions document [RD.177]</i> .	CF	This is not used in SST_cci files.
flag_meanings	string	Space-separated list of text descriptions. Words within a phrase should be connected with underscores. Used only for flag and "sources_of_xxx" variables.	CF	See specifications in Section 4.
flag_values	Array of the data type of variable (Note that this is listed incorrectly as 'string' in [AD.1])	Comma-separated array of valid, mutually exclusive variable values (required when the bit field contains enumerated values). Used only for flag and "sources_of_xxx" variables.	CF	See specifications in Section 4.
flag_masks	Array of the data type of variable (Note that this is listed incorrectly as 'string' in [AD.1])	Array of valid variable masks (required when the bit field contains independent Boolean conditions). Used only for flag and "sources_of_xxx" variables.	CF	See specifications in Section 4.

3.5 Unique product identifiers

Some of the attributes described in Table 6 and Table 8 contain identifiers for the data that have been used to create the file. Section 7.9 of [AD.1] describes these strings. For completeness a brief summary can be found below. In Table 9 all the source data to be used by the project are listed together with the unique text strings that shall be used by to refer to these in the SST_cci files. The entries to the table are based on the information in [AD.1] and [RD.172].

The format of the text is:

<Product String>-<RDAC>-<Processing Level>-<Additional Segregator>-v<Product Version>

The definition of these elements can be found in Table 3, with the exception of the following points:

- <Product Version> - this specifies the version of the dataset and should have the form x.y, where x is the major and y the minor version number.
- <Additional Segregator> - for wind speed data, <Additional Segregator> should be set to WSP, and to ICE for sea ice fraction.
- If there is no predefined GHRSSST RDAC for a data source an appropriate acronym can be substituted instead.

[AD.1] includes a number of predefined text strings. Some existing text strings predate the format described above and hence do not conform.

Table 9. Table of source data and unique product identifiers. The unique text string for SLSTR data should be checked against the GHRSSST specification before use.

Source data	Unique text string	Comment
ATSR1 Level 1	ATSR1-ESA-L1-v2.0	
ATSR2 Level 1	ATSR2-ESA-L1-v2.0	
AATSR Level 1	AATSR-ESA-L1-v2.0	
AVHRR GAC Level 1	AVHRR<X>_G-NOAA-L1-v<Y>	<X> is the satellite number; <Y> is the format version number in form x.y where x is the major format version (1, 2, 3, 4 or 5) and y is used to denote format changes where the major format number is unchanged (possible values are 0, 1 and 2).
AVHRR MetOp Level 1	AVHRRMTA-OSISAF-L1B-vA1.0p1.0	Version number format is non-standard.
SLSTR Level 1	S3A_SLSTR-ESA-L1-v<x.y>	TBC
NWP wind speed	ERA_INTERIM-ECMWF-WSP-v1.0	
OSI SAF sea ice fractions	EUMETSAT_OSI-SAF-ICE-v1.1	This is an adaptation of the source field used in current OSTIA files
Land mask	NAVOCEANO_landmask_v1.0	
Lake mask	ARCLake_lakemask	
Data files created by SST_cci	<Product String>-ESACCI-<Processing Level>-v<Version Number>	See text for definition of these fields.

4. PRODUCT SPECIFICATION

This section contains detailed specifications for the files to be produced by the SST_cci project. The structure of the files is fully specified, including the geophysical variables, the metadata and aspects of the data such as the grid.

In addition to the tables in this section, Network Common Data Form Language (CDL) format listing of the headers of example L2P, L3U and L4 NetCDF files are provided in Appendix A. These have been tested using the CF compliance checker at <http://puma.nerc.ac.uk/cgi-bin/cf-checker.pl> to confirm that they comply with the CF conventions version 1.5.

4.1 Format of the product description tables

Tables are used to describe the detailed structure of the data files in the following subsections. An example of the layout of the tables is shown in Table 10. The points to note are:

- Colours are used to show the start and end of sections of the file structure.
- The column headed 'T' contains the storage type of the variables and attributes; the abbreviations used are explained in Table 11.
- The column headed 'D' contains the dimensionality of the variables. Dimensions are ordered as in the network Common Data form Language (CDL) definition i.e. with the fastest changing dimension last.
- The 'Bytes' column indicates the storage requirement for each item.

Table 10. Example of a table showing file structure. Colours are used to denote the beginning and end of sections of the structure.

Element name	Description	Range/value	Unit	T	D	Bytes
Dataset	Begin of dataset					
dimensions	Begin of dimensions					
dimension1						
dimension2						
dimensionN						
dimensions	End of dimensions					
variables	Begin of variables					
variable	Begin of variable					
variable_name						
attributes	Begin of attributes					
attribute1						
attribute2						
attributeN						
attributes	End of attributes					
variable	End of variable					

Element name	Description	Range/value	Unit	T	D	Bytes
More variable definitions as required						
variables	End of variables					
attributes	Begin of global attributes					
Global attributes that have values that are consistent between data levels are defined in Table 13.						
global_attribute1						
global_attribute2						
global_attributeN						
attributes	End of global attributes					
dataset	End of dataset					

4.2 Storage type definitions

Storage types and abbreviations used to refer to them are listed in Table 11. Storage types are limited to those available in the classic NetCDF format. For information the names given to these types in [AD.1] are also listed.

Note that although the abbreviation 'st' is used to indicate text strings, in the NetCDF classic data model there is actually no 'string' storage type [RD.167]. This type of data is instead stored as an array of characters.

Table 11. Abbreviations used for storage types.

Type	Description	Name used in [AD.1]	Comment
sc	8-bit signed integer	byte	The NetCDF data type names match those used in [AD.1] for these types.
ss	16-bit signed integer	short	
sl	32-bit signed integer	int (or long)	
fl	32-bit floating point	float	
db	64-bit floating point	double	
st	Character array	string	To be stored in the NetCDF file as an array of characters (8-bit unsigned integers)

4.3 L2P data

4.3.1 Description

L2P data files contain swath based SST data. In the context of the SST_cci project, they will be produced from AVHRR Global Area Coverage (GAC) data.

Each file shall contain a single orbit of data. The pixel geometry used is the sub-sampled geometry of GAC with its variable resolution across the swath.

L2P files shall include confidence information and single sensor error statistics (SSES) (bias and standard deviation).

Multiple different types of retrieval are possible using different views and channels. The L2P files shall contain the best available retrievals. No specific user requirement has been identified for files containing the other retrievals. However, in case such a requirement becomes apparent a method of storing these data is defined in Appendix B.

4.3.2 Summary of contents

The files produced by SST_cci will meet the specification of the 'L2Pcore' files defined in [AD.1]. The SSES mean bias traditionally provides a bias correction to apply to the data. This is not appropriate in the case of SST_cci products, so SSES bias will be populated with zeros. SSES standard deviations will contain the total uncertainty of each data point of the best available skin/subskin SSTs. One variable from the list of auxiliary fields (L2Paux), wind speed, is also included within the SST_cci L2P files. In addition, six experimental variables are included:

- SST adjusted for time and depth,
- large scale correlated uncertainty (highly correlated between pixels over long spatiotemporal scales),
- synoptically correlated uncertainty (highly correlated between pixels over synoptic scales),
- uncorrelated uncertainty (uncorrelated or weakly correlated between pixels), and,
- uncertainty in time and depth adjustment (a systematic uncertainty),
- total uncertainty for the SST adjusted for time and depth.

Information about how to combine the uncertainties is provided in the comment attributes within the files and correlation length and time scales are provided for the synoptically correlated components. CF standard name modifiers do not exist for individual uncertainty components and so no standard names are defined for them. It is suggested that these should be proposed in the future if it becomes clear that there is a general need for uncertainties that are decomposed in this way.

A summary of the data fields is given in Table 12.

Table 12. Variables to be included within the L2P data files; some of the name and description information in this table is reproduced from Section 9.1 of [AD.1]. The shading (see Table 7) and the OCC column indicate the number of occurrences of each variable. These are either mandatory (1, blue shading) or optional (0..1, violet shading).

Name	Description	OCC
lat	Coordinate variable; central latitude of each spatial point of the data arrays	1
lon	Coordinate variable; central longitude of each spatial point of the data arrays	1
time	Coordinate variable; time of each temporal point of the data arrays; the start time of the granule	1
sea_surface_temperature	Best available skin/subskin SST retrievals (unadjusted for time and depth); fill values to be provided where there is land, cloud or total sea ice cover	1

Name	Description	OCC
sst_dtime	Time differences of SST retrievals from the base time in the time coordinate variable	1
sses_bias	SST single sensor error statistic measurement bias estimate (to contain zeroes)	1
sses_standard_deviation	SST single sensor error statistic measurement standard deviation estimate (to contain the total of the large scale correlated, synoptically correlated and uncorrelated uncertainties)	1
l2p_flags	Flags specific to each L2P dataset that help users interpret data	1
quality_level	A quality level for each measurement	1
wind_speed	An estimate of surface wind speed (from the ERA-Interim reanalysis)	1
sea_surface_temperature_depth	Sea surface temperature at standard depth and time (adjusted version of the sea_surface_temperature variable)	1
sst_depth_dtime	Time differences from the base time in the time coordinate variable for the SSTs adjusted to standard depth and time.	1
large_scale_correlated_uncertainty	Systematic uncertainty that is highly correlated between pixels over large scales	1
synoptically_correlated_uncertainty	Systematic uncertainty that is highly correlated between pixels over synoptic scales only	1
uncorrelated_uncertainty	Non-systematic uncertainty (uncorrelated or weakly correlated between pixels)	1
adjustment_uncertainty	Uncertainty in adjustment to standard depth and time (correlated between pixels over synoptic scales)	1
sst_depth_total_uncertainty	To contain the total of the large scale correlated, synoptically correlated, uncorrelated and adjustment uncertainties.	1

The contents of these files meet the following user requirements:

- The inclusion of wind speed meets user requirement SST_CCI-URD-QUF-70 [RD.171].
- The SST that is adjusted for time and depth will correspond to 10.30 am or pm and a depth of 20 cm. This is the (joint) second most popular choice of depth with users according to requirement SST_CCI-UR-QUF-46 [RD.171]. Note that the most popular choice (skin SST) corresponds to the SST viewed by infrared instruments and so will also be included in the files for data from that type of sensor.
- The separate uncertainty components and correlation scales provide information to the user about the correlation structure of the uncertainties (requirement SST_CCI-UR-QUF-57, [RD.171]).

- When the uncertainty components are combined they give an estimate of total uncertainty (requirement SST_CCI-UR-QUF-53, [RD.171]).
- The files also include a simple quality flag (SST_CCI-URD-QUF-58, [RD.171]).

4.3.3 Detailed contents

The detailed contents of the L2P products are defined in Table 13 and Table 14. The former is used to specify the global attributes that have common values in files for all data levels. Together the tables specify the structure of the NetCDF files. Examples of what the attributes might contain in a real file can be found in Appendix A.

Table 13. Global attributes common to all data files. Contents were taken from [AD.1] Table 8-1 and adapted for the SST_cci data files.

Element name	Description	Range/ value	Unit	T	D	Bytes
attributes	Beginning of global attributes					
Conventions	The version of the netCDF conventions followed.	CF-1.5, Unidata Observation Dataset v1.0		st		
title	Insert satellite sensor name and data level into the string to create a descriptive name for the data.	ESA SST CCI <Product String> <Data Level> product		st		
summary	Insert satellite sensor name, data level and algorithm name/brief description.	<Product String> <Data Level> product from the ESA SST CCI project, produced using <algorithm name>.		st		
references	Published or web-based references.			st		
institution	RDAC code.	ESACCI		st		

Element name	Description	Range/ value	Unit	T	D	Bytes
history	To contain a history of applications that have been used to process the data. For example this can be used to record the version of the retrieval algorithm applied to the data, the identity of intermediate files used in the processing etc.			st		
license	The GHRSSST standard data license.	GHRSSST protocol describes data use as free and open		st		
id	Unique text string for this product. See Section 3.5.			st		
naming_authority	Defined in [RD.87] as org.ghrsst.	org.ghrsst		st		
product_version	Product version (see Section 3.5).			st		
uuid	Universally Unique Identifier. For example see http://www.ossfp.org/pkg/lib/uuid/ [RD.169]			st		
tracking_id	Identical to uuid (i.e. this shall contain the same value as uuid)			st		
gds_version_id	Version of [GDS] that files are based on.	2.0		st		
netcdf_version_id	NetCDF library.	Format examples: 4.1.1 or 4.2.1.1 of Oct 19 2012 14:25:16		st		
date_created	File creation	Format: yyyymmddThh mmssZ		st		

Element name	Description	Range/ value	Unit	T	D	Bytes
file_quality_level	Select from: 0 = unknown quality 1 = extremely suspect (frequent problems, e.g. with known satellite problems) 2 = suspect (occasional problems, e.g. after launch) 3 = excellent (no known problems)	[0, 3]		sl		4
spatial_resolution	A string describing the approximate resolution of the product. For example, "1.1km at nadir". See Table 1 for values to use.	[1 km at nadir, 0.05°]		st		
start_time	Date and time of the first measurement in the data file.	Format: yyyyymmddThhmmssZ		st		
time_coverage_start	Identical to start_time.	Format: yyyyymmddThhmmssZ		st		
stop_time	Date and time of the last measurement in the data file in the form.	Format: yyyyymmddThhmmssZ		st		
time_coverage_end	Identical to stop_time.	Format: yyyyymmddThhmmssZ		st		
time_coverage_duration	Difference between time_coverage_end and time_coverage_start in the form PdDThHmMsS, replacing the lower case letters with the appropriate numbers.	Format: PdDThHmMsS (see description column for more details) or other ISO 8601 compliant format		st		

Element name	Description	Range/ value	Unit	T	D	Bytes
time_coverage_resolution	Temporal resolution of data in the file i.e. the orbit repeat period or the frequency of L3/L4 data.	Format: PdDThHmMs S (see description of time_coverage_duration for more details)		st		
source	Comma separated list of all source data present in this file. List SST sources first, followed by auxiliary sources. See Section 3.5 for source data names.			st		
platform	Satellite(s) used to create this data file. Select from the entries found in Table 5			st		
sensor	Satellite(s) used to create this data file. Select from the entries found in Table 5			st		
Metadata_Conventions		Unidata Dataset Discovery v1.0		st		
metadata_link	Link to collection metadata record at archive.	http://www.esa-cci.org		st		
keywords	Standard words that describe the data, taken from the source specified in keywords_vocabulary.	Oceans > Ocean Temperature > Sea Surface Temperature		st		
keywords_vocabulary	Defines the source of the text in the keywords attribute; [RD.176].	NASA Global Change Master Directory (GCMD) Science Keywords		st		

Element name	Description	Range/ value	Unit	T	D	Bytes
standard_name_vocabulary	Defines the source of the standard names for the variables; [RD.177].	NetCDF Climate and Forecast (CF) Metadata Convention		st		
geospatial_lat_units	Units of the latitudinal resolution.	degrees_north		st		
geospatial_lat_resolution	Latitude resolution in units matching geospatial_lat_units. See values in Table 1.	[0.009, 0.05]		fl		4
geospatial_lon_units	Units of the longitudinal resolution.	degrees_east		st		
geospatial_lon_resolution	Longitude resolution in units matching geospatial_lon_units. See values in Table 1.	[0.009, 0.05]		fl		4
geospatial_vertical_min	Depth of the deepest SST in the file (negative value as the direction is downwards) – see Table 4	-10 ⁻⁶ or -0.001 or -0.2 or -10		fl		4
geospatial_vertical_max	Depth of the shallowest SST in the file (negative value as the direction is downwards) – see Table 4	-10 ⁻⁶ or -0.001 or -0.2 or -10		fl		4
acknowledgment	Information about funding source and how to cite the use of these data.	Funded by ESA		st		
creator_name	Description of data creators.	ESA SST CCI		st		
creator_email		science.leader@esa-sst-cci.org		st		
creator_url		http://www.esa-sst-cci.org/		st		

Element name	Description	Range/ value	Unit	T	D	Bytes
creator_processing_institution	Contains additional information about the institution that created the product within the CCI project. Insert institution name (for example 'the University of Edinburgh' or 'the Met Office') to indicate where the data were produced.	These data were produced at <institution> as part of the ESA SST CCI project.		st		
project	The name of the project.	Climate Change Initiative – European Space Agency		st		
publisher_name	Information about the data publisher.	ESACCI		st		
publisher_url		http://www.esa-sst-cci.org		st		
publisher_email		science.leader@esa-SST-cci.org		st		
product_specification_version	Version of this document used to define the file.	The filename of the PSD (without extension)		st		
attributes	End of global attributes					

Table 14. Detailed specification of the SST_cci L2P files.

Element name	Description	Range/value	Unit	T	D	Bytes
dataset	Begin of dataset					
dimensions	Begin of dimensions					
ni	Across track dimension.			sl	1	4
nj	Along track dimension.			sl	1	4
time	Time dimension; must be 1 for L2P data.	1		sl	1	4
dimensions	End of dimensions					

Element name	Description	Range/value	Unit	T	D	Bytes
variables	Begin of variables					
variable	Begin of variable					
lat	Latitude coordinates.	[-90, 90]		fl	nj, ni	nj × ni × 4
attributes	Begin of attributes					
units	Text description of the units.	degrees_north		st		
long_name	A free-text descriptive variable name.	Latitude coordinates		st		
standard_name	Unique descriptive name for data.	latitude		st		
valid_min	Minimum valid value for this variable.	-90.0		fl		4
valid_max	Maximum valid value for this variable.	90.0		fl		4
reference_datum	Information about the coordinates.	geographical coordinates, WGS84 projection		st		
comment	Miscellaneous information about the data or the methods used to produce it. (Optional.)			st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
lon	Longitude coordinates.	[-180, 180]		fl	nj, ni	nj × ni × 4
attributes	Begin of attributes					
units	Text description of the units.	degrees_east		st		
long_name	A free-text descriptive variable name.	Longitude coordinates		st		
standard_name	Unique descriptive name for data.	longitude		st		

Element name	Description	Range/value	Unit	T	D	Bytes
valid_min	Minimum valid value for this variable.	-180.0		fl		4
valid_max	Maximum valid value for this variable.	180.0		fl		4
reference_datum	Information about the coordinates.	geographical coordinates, WGS84 projection		st		
comment	Miscellaneous information about the data or the methods used to produce it. (Optional.)			st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
time	Time coordinate ; the start time of the granule	[0, 2 ³¹]		sl	time	time x 4
attributes	Begin of attributes					
units	Text description of the units.	seconds since 1981-01-01 00:00:00		st		
long_name	A free-text descriptive variable name.	reference time of sst file		st		
standard_name	Unique descriptive name for data.	time		st		
calendar	Defines the calendar used to define the times.	gregorian		st		
comment	Miscellaneous information about the data or the methods used to produce it. (Optional.)			st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					

Element name	Description	Range/value	Unit	T	D	Bytes
sea_surface_temperature	Best available sea surface temperature retrievals; to contain fill values if land, cloud or full sea ice cover.	[271.15, 323.15] (after scaling)	K	ss	time, nj, ni	time x nj x ni x 2
attributes	Beginning of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-32768		ss		2
units	Text description of the units.	kelvin		st		
scale_factor	To be multiplied by the variable to recover the original value.	0.01		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	273.15		fl		4
long_name	A free-text descriptive variable name. First option is for infrared data, second for passive microwave data; see also Table 4.	sea surface skin temperature or sea surface subskin temperature		st		
valid_min	Minimum valid value for this variable once they are packed (in storage type).	-200		ss		2
valid_max	Maximum valid value for this variable once they are packed (in storage type).	5000		ss		2

Element name	Description	Range/value	Unit	T	D	Bytes
standard_name	Unique descriptive name for data. Choose appropriate value from Table 4.	sea_surface_skin_temperature or sea_surface_subskin_temperature		st		
comment	Miscellaneous information about the data or the methods used to produce it. Includes method of calculating total uncertainty.	Temperature of the skin of the ocean; total uncertainty = $\sqrt{\text{large_scale_correlated_uncertainty}^2 + \text{synoptically_correlated_uncertainty}^2 + \text{uncorrelated_uncertainty}^2}$		st		
source	See Table 9.			st		
references	Published or web-based references that describe the data or methods used to produce it.			st		
depth	Effective depth of the sea_surface_temperature variable. Choose appropriate value from Table 4.	10 micrometres or 1 millimetre		st		
coordinates	Identifies coordinate variables.	lon lat		st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
sst_dtime	Time differences of SST retrievals from the reference time.	[-32767, 32767]	secs	ss	time, nj, ni	time × nj × ni × 2
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-32768		ss		2

Element name	Description	Range/value	Unit	T	D	Bytes
units	Text description of the units.	seconds		st		
scale_factor	To be multiplied by the variable to recover the original value.	1.0		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	0.0		fl		4
long_name	A free-text descriptive variable name.	time difference from reference time		st		
valid_min	Minimum valid value for this variable once they are packed (in storage type).	-32767		ss		2
valid_max	Maximum valid value for this variable once they are packed (in storage type).	32767		ss		2
comment	Miscellaneous information about the data or the methods used to produce it.	time plus sst_dtime gives seconds after 1981-01-01 00:00:00		st		
coordinates	Identifies coordinate variables.	lon lat		st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
sses_bias	Single sensor error statistic bias error.	Zero for all SST values	K	sc	time, nj, ni	time x nj x ni
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-128		sc		1

Element name	Description	Range/value	Unit	T	D	Bytes
units	Text description of the units.	kelvin		st		
scale_factor	To be multiplied by the variable to recover the original value.	0.01		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	0.0		fl		4
long_name	A free-text descriptive variable name.	SSES bias estimate		st		
valid_min	Minimum valid value for this variable once they are packed (in storage type).	-127		sc		1
valid_max	Maximum valid value for this variable once they are packed (in storage type).	127		sc		1
comment	Miscellaneous information about the data or the methods used to produce it.			st		
coordinates	Identifies coordinate variables.	lon lat		st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
sses_standard_deviation	Single sensor error statistic standard deviation.	Total uncertainty from combining uncertainties as described in the comment attribute to the sea_surface_temperature variable	K	sc	time, nj, ni	time × nj × ni
attributes	Begin of attributes					

Element name	Description	Range/value	Unit	T	D	Bytes
_FillValue	A value used to indicate array elements containing no valid data.	-128		sc		1
units	Text description of the units.	kelvin		st		
scale_factor	To be multiplied by the variable to recover the original value.	0.01		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	1.27		fl		4
long_name	A free-text descriptive variable name.	SSES standard deviation		st		
valid_min	Minimum valid value for this variable once they are packed (in storage type).	-127		sc		1
valid_max	Maximum valid value for this variable once they are packed (in storage type).	127		sc		1
comment	Miscellaneous information about the data or the methods used to produce it. Only include the text ' adjustment_ uncertainty' if this is to be included in the file (i.e. do not include if it is a demonstration ECV file).	Uncertainty data are also contained in the variables large_scale_correlated_ uncertainty, synoptically_correlated_ uncertainty, uncorrelated_ uncertainty and adjustment_ uncertainty.		st		
coordinates	Identifies coordinate variables.	lon lat		st		
attributes	End of attributes					
variable	End of variable					

Element name	Description	Range/value	Unit	T	D	Bytes
variable	Begin of variable					
l2p_flags	Specifies type of input data and provide flags and information to the user. See Table 15.	[0, 255] (this may change if extra flags are defined)		ss	time x nj x ni	time x nj x ni x 2
attributes	Begin of attributes					
valid_min	Minimum valid value for this variable once they are packed (in storage type).	0		sc		1
valid_max	Maximum valid value for this variable once they are packed (in storage type).	255		sc		1
long_name	A free-text descriptive variable name.	L2P flags		st		
comment	Miscellaneous information about the data or the methods used to produce them.	These flags are important to properly use the data		st		
coordinates	Identifies coordinate variables.	lon lat		st		
flag_meanings	Meaning attached to each flag value.	microwave land ice lake river spare views channels		st		
flag_masks	Bit masks corresponding to the flags described in flag_meanings.	1, 2, 4, 8, 16, 32, 64, 128		ss		2
attributes	End of attributes					
variable	Begin of variable					
quality_level	Indicator of the quality of each SST.	[0, 5]		sc	time x nj x ni	time x nj x ni
attributes	Begin of attributes					

Element name	Description	Range/value	Unit	T	D	Bytes
_FillValue	A value used to indicate array elements containing no valid data.	-128		sc		1
valid_min	Minimum valid value for this variable once they are packed (in storage type).	0		sc		1
valid_max	Maximum valid value for this variable once they are packed (in storage type).	5		sc		1
long_name	A free-text descriptive variable name.	quality level of SST pixel		st		
comment	Miscellaneous information about the data or the methods used to produce it.	These are the overall quality indicators and are those used for all GHRSSST SSTs		st		
coordinates	Identifies coordinate variables.	lon lat		st		
flag_meanings	Meaning attached to each flag value.	no_data bad_data worst_quality low_quality acceptable_q uality best_quality		st		
flag_values	Values that correspond to the flags described in flag_meanings. The b after the numbers is to indicate that they are single byte numbers.	0b, 1b, 2b, 3b, 4b, 5b		sc		1
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					

Element name	Description	Range/value	Unit	T	D	Bytes
wind_speed	Surface wind speed at 10 m height. Resolution should be no less than 1 m s ⁻¹ and has been defined here to be 0.1 m s ⁻¹ . The range of wind speeds that can be represented is 0 – 25.4 m s ⁻¹ ; values higher than this should be set to the maximum.	[0, 25.4] (after scaling)	ms ⁻¹	sc	time, nj, ni	time x nj x ni
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-128		sc		1
units	Text description of the units.	m s-1		st		
scale_factor	To be multiplied by the variable to recover the original value.	0.1		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	12.7		fl		4
long_name	A free-text descriptive variable name.	10m wind speed		st		
valid_min	Minimum valid value for this variable once they are packed (in storage type).	-127		sc		1
valid_max	Maximum valid value for this variable once they are packed (in storage type).	127		sc		1
standard_name	Unique descriptive name for data.	wind_speed		st		

Element name	Description	Range/value	Unit	T	D	Bytes
comment	Miscellaneous information about the data or the methods used to produce it.	Wind speeds sourced from ECMWF ERA Interim reanalysis; wind speeds greater than 25.4 m/s are set to 25.4.		st		
source	See Table 9.			st		
references	Published or web-based references that describe the data or methods used to produce it.			st		
coordinates	Identifies coordinate variables.	lon lat		st		
height	Height that the wind values correspond to.	10 m		st		
time_offset	Difference in hours between the SST reference time and the time of the wind speed value.			fl		4
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
sea_surface_temperature _depth	SSTs adjusted to 10.30 local time (either morning or evening, whichever is closest) and 20 cm.	[271.15, 323.15] (after scaling)	K	ss	time, nj, ni	time x nj x ni x 2
attributes	Beginning of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-32768		ss		2
units	Text description of the units.	kelvin		st		

Element name	Description	Range/value	Unit	T	D	Bytes
scale_factor	To be multiplied by the variable to recover the original value.	0.01		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	273.15		fl		4
long_name	A free-text descriptive variable name.	sea surface temperature at 0.2 m		st		
valid_min	Minimum valid value for this variable once they are packed (in storage type).	-200		ss		2
valid_max	Maximum valid value for this variable once they are packed (in storage type).	5000		ss		2
standard_name	Unique descriptive name for data.	sea_water_temperature		st		
comment	Miscellaneous information about the data or the methods used to produce it. Includes method of calculating total uncertainty.	Temperature of the ocean at 20 cm depth; total uncertainty = $\sqrt{\text{large_scale_correlated_uncertainty}^2 + \text{synoptically_correlated_uncertainty}^2 + \text{uncorrelated_uncertainty}^2 + \text{adjustment_uncertainty}^2}$		st		
source	See Table 9.			st		
references	Published or web-based references that describe the data or methods used to produce it.			st		
depth	Effective depth of the SST data.	0.2 metre		st		

Element name	Description	Range/value	Unit	T	D	Bytes
coordinates	Identifies coordinate variables.	lon lat		st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
sst_depth_dtime	Time differences of SST depth from the reference time.	[-32767, 32767]	secs	ss	time, nj, ni	time × nj × ni × 2
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-32768		ss		2
units	Text description of the units.	seconds		st		
scale_factor	To be multiplied by the variable to recover the original value.	1.0		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	0.0		fl		4
long_name	A free-text descriptive variable name.	time difference from reference time		st		
valid_min	Minimum valid value for this variable once they are packed (in storage type).	-32767		ss		2
valid_max	Maximum valid value for this variable once they are packed (in storage type).	32767		ss		2
comment	Miscellaneous information about the data or the methods used to produce it.	time plus sst_dtime gives seconds after 1981-01-01 00:00:00		st		

Element name	Description	Range/value	Unit	T	D	Bytes
coordinates	Identifies coordinate variables.	lon lat		st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
large_scale_correlated_uncertainty	Component of uncertainty that is highly correlated between SST retrievals separated by larger than sub-synoptic spatio-temporal scales.	[0, 50] (after scaling)	K	ss	time x nj x ni	time x nj x ni x 2
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-32768		ss		2
units	Text description of the units.	kelvin		st		
scale_factor	To be multiplied by the variable to recover the original value.	0.01		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	0.0		fl		4
long_name	A free-text descriptive variable name.	Uncertainty from errors likely to be correlated over large scales		st		
valid_min	Minimum valid value for this variable once they are packed (in storage type).	0		ss		2
valid_max	Maximum valid value for this variable once they are packed (in storage type).	5000		ss		2

Element name	Description	Range/value	Unit	T	D	Bytes
comment	Miscellaneous information about the data or the methods used to produce it.	Component of uncertainty that is correlated over large scales; can be combined with other uncertainty estimates to form a total uncertainty		st		
references	Published or web-based references that describe the data or methods used to produce it.			st		
coordinates	Identifies coordinate variables.	lon lat		st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
synoptically_correlated_uncertainty	Component of uncertainty that is highly correlated between SST retrievals separated by sub-synoptic spatio-temporal scales.	[0, 50] (after scaling)	K	ss	time x nj x ni	time x nj x ni x 2
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-32768		ss		2
units	Text description of the units.	kelvin		st		
scale_factor	To be multiplied by the variable to recover the original value.	0.01		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	0.0		fl		4

Element name	Description	Range/value	Unit	T	D	Bytes
long_name	A free-text descriptive variable name.	Uncertainty from errors like to be correlated over synoptic scales		st		
valid_min	Minimum valid value for this variable once they are packed (in storage type).	0		ss		2
valid_max	Maximum valid value for this variable once they are packed (in storage type).	5000		ss		2
comment	Miscellaneous information about the data or the methods used to produce it.	Component of uncertainty that is correlated over synoptic scales; can be combined with other uncertainty estimates to form a total uncertainty		st		
references	Published or web-based references that describe the data or methods used to produce it.			st		
coordinates	Identifies coordinate variables.	lon lat		st		
correlation_length_scale	Estimate of the correlation length scale of the uncertainties.	100 km		st		
correlation_time_scale	Estimate of the correlation time scale of the uncertainties	1 day		st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					

Element name	Description	Range/value	Unit	T	D	Bytes
uncorrelated_uncertainty	Component of uncertainty that is uncorrelated between SST retrievals separated by any spatio-temporal scales.	[0, 50] (after scaling)	K	ss	time x nj x ni	time x nj x ni x 2
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-32768		ss		2
units	Text description of the units.	kelvin		st		
scale_factor	To be multiplied by the variable to recover the original value.	0.01		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	0.0		fl		4
long_name	A free-text descriptive variable name.	Uncertainty from errors likely to be uncorrelated between SSTs		st		
valid_min	Minimum valid value for this variable once they are packed (in storage type).	0		ss		2
valid_max	Maximum valid value for this variable once they are packed (in storage type).	5000		ss		2

Element name	Description	Range/value	Unit	T	D	Bytes
comment	Miscellaneous information about the data or the methods used to produce it.	Component of uncertainty that is uncorrelated between SSTs; can be combined with other uncertainty estimates to form a total uncertainty		st		
references	Published or web-based references that describe the data or methods used to produce it.			st		
coordinates	Identifies coordinate variables.	lon lat		st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
adjustment_uncertainty	Uncertainty in adjustment for depth and time, correlated between SST estimates separated by sub-synoptic spatio-temporal scales.	[0, 50] (after scaling)	K	ss	time x nj x ni	time x nj x ni x 2
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-32768		ss		2
units	Text description of the units.	kelvin		st		
scale_factor	To be multiplied by the variable to recover the original value.	0.01		fl		4

Element name	Description	Range/value	Unit	T	D	Bytes
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	0.0		fl		4
long_name	A free-text descriptive variable name.	Time and depth adjustment uncertainty		st		
valid_min	Minimum valid value for this variable once they are packed (in storage type).	0		ss		2
valid_max	Maximum valid value for this variable once they are packed (in storage type).	5000		ss		2
comment	Miscellaneous information about the data or the methods used to produce it.	Adjustment uncertainty; can be combined with other uncertainty estimates to form a total uncertainty		st		
references	Published or web-based references that describe the data or methods used to produce it.			st		
coordinates	Identifies coordinate variables.	lon lat		st		
correlation_length_scale	Estimate of the correlation length scale of the uncertainties.	100 km		st		
correlation_time_scale	Estimate of the correlation time scale of the uncertainties	1 day		st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					

Element name	Description	Range/value	Unit	T	D	Bytes
sst_depth_total_uncertainty	Combination of all uncertainty components.	Total uncertainty from combining uncertainties as described in the comment attribute to the sea_surface_temperature_depth variable	K	ss	time, nj, ni	time x nj x ni x 2
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-32768		ss		2
units	Text description of the units.	kelvin		st		
scale_factor	To be multiplied by the variable to recover the original value.	0.01		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	0.00		fl		4
long_name	A free-text descriptive variable name.	Total uncertainty in sea_surface_temperature_depth		st		
valid_min	Minimum valid value for this variable once they are packed (in storage type).	0		ss		2
valid_max	Maximum valid value for this variable once they are packed (in storage type).	5000		ss		2
comment	Miscellaneous information about the data or the methods used to	Total uncertainty in each sea_surface_temperature_		st		

Element name	Description	Range/value	Unit	T	D	Bytes
	produce it.	depth data point				
coordinates	Identifies coordinate variables.	lon lat		st		
attributes	End of attributes					
variable	End of variable					
variables	End of variables					
attributes	Begin of global attributes					
Global attributes as described in Table 6.						
comment	To contain any miscellaneous information about the data and methods. Delete the 'and sea_surface_temperature_depth' part of the second sentence if the sea_surface_temperature_depth variable is not included in the file.	For information about uncertainty estimates see the comment attributes to the sea_surface_temperature and sea_surface_temperature_depth variables.		st		
northernmost_latitude	Decimal degrees north, range -90 to +90. May vary from orbit to orbit.	[-90, 90]		fl		4
geospatial_lat_max	Identical to northernmost_latitude	[-90, 90]		fl		4
southernmost_latitude	Decimal degrees north, range -90 to +90. May vary from orbit to orbit.	[-90, 90]		fl		4
geospatial_lat_min	Identical to southernmost_latitude	[-90, 90]		fl		4
easternmost_longitude	Decimal degrees east, range -180 to +180. May vary from orbit to orbit.	[-180, 180]		fl		4
geospatial_lon_max	Identical to easternmost_longitude	[-180, 180]		fl		4

Element name	Description	Range/value	Unit	T	D	Bytes
westernmost_longitude	Decimal degrees east, range -180 to +180. May vary from orbit to orbit.	[-180, 180]		fl		4
geospatial_lon_min	Identical to westernmost_longitude	[-180, 180]		fl		4
processing_level	Data level.	L2P		st		
cdm_data_type	“swath” or “grid”	swath		st		
attributes	End of global attributes					
dataset	End of dataset					

Table 15. Meaning of each bit in a l2p_flags field. Reformatted version of Table 9-19 of [AD.1] with additional entries denoted by italics. Bits 6-15 are available for use by the SST_cci project; two of these are used to describe the retrieval.

Bit	Description	Meaning if 0	Meaning if 1
0	Type of sensor	Infrared	Microwave
1	Ocean or land indicator	Over the ocean	Over land
2	Sea ice indicator	Not sea ice contaminated	Sea ice contaminated
3	Lake indicator (if known)	Not over lake	Over lake
4	River indicator (if known)	Not over river	Over river
5	Reserved for future use		
6	<i>Number of satellite views used in retrieval (or, if no retrieval possible, what it would have been if a retrieval had been performed))</i>	<i>Single (nadir only) view</i>	<i>Dual view</i>
7	<i>Number of channels used in retrieval (or, if no retrieval possible, what it would have been if a retrieval had been performed)</i>	<i>Two channels</i>	<i>Three channels</i>
8-15	<i>Available for use by SST_cci</i>		

Table 16. Quality level indicators for SST data. To simplify the flags SST_cci shall use only values of 0, 1 and 5. Reproduced from Section 9.18 of [AD.1] with shading added to indicate flags that will not be used by SST_cci.

Value	Meaning
0	Missing data
1	These data are not valid and should not be used
2	Data are of the lowest quality that still could be usable
3	Low quality data
4	Acceptable quality data
5	Best quality data

4.4 L3U data

4.4.1 Description

Level 3 data files contain gridded SST data (but not analysis data, i.e. they do not include values where no retrieval could be made). In the context of the data to be produced by SST_cci these will be uncollated (L3U) data files – these contain a single orbit of data from a single sensor (i.e. a gridded version of L2P data). L3U files will be produced for ATSR series data.

The grid used shall be a regular 0.05° latitude-longitude grid (SST_CCI-UR-QUF-36, [RD.171]). As with L2P files, the level 3 files shall contain the best available SST data. See Appendix B for specifications of files that could be used to hold data obtained using other retrievals should a user requirement arise. L3C files – daily collations of data from a sensor – can also be produced following the specification described in this document.

The contents of the files are very similar to those of L2P files and hence meet the same user requirements described in Section 4.3.2.

4.4.2 Summary of contents

The files are almost identical to the L2P files. They differ in the specification of the grid and in some of the attributes. Note that the time dimension in level 3 (and 4) files is set to be unlimited. In practice only one time field is contained in each file, but this allows utilities that operate on the files to concatenate the data [AD.1].

Table 17. Fields within a L3 data file; some of the name and description information in this table is reproduced from Section 9.1 of [AD.1]. The shading (see Table 7) and the OCC column indicate the number of occurrences of each variable. These are either mandatory (1, blue shading) or optional (0..1, violet shading).

Name	Description	OCC
lat	Vector specifying the central latitude of each spatial point in the data grid	1
lat_bnds	Latitude bounds of each grid cell; included from [RD.166].	1
lon	Vector specifying the central longitude of each spatial point in the data grid	1
lon_bnds	Longitude bounds of each grid cell; included from [RD.166].	1
time	Vector specifying the reference time of each temporal point in the data grid; this is either the start time of the L3U granule or the mid-point of the collation window for L3C (the midpoint of the day the data represent)	1
time_bnds	Time bounds of each time point (either start and end times of the data collection for L3U or start and end of the day for L3C); included from [RD.166]	1
sea_surface_temperature	Best available sea surface temperature retrievals (unadjusted for time and	1

	depth); fill values to be provided where there is land, cloud or total sea ice cover	
sst_dtime	Time differences of SST retrievals from the reference time	1
sses_bias	SST single sensor error statistic measurement bias estimate (to contain zeroes)	1
sses_standard_deviation	SST single sensor error statistic measurement standard deviation estimate (to contain the total of the large scale correlated, synoptically correlated and uncorrelated uncertainties)	1
l2p_flags	Flags that help users interpret data	1
quality_level	A quality level for each SST	1
wind_speed	An estimate of surface wind speed (from the ERA-Interim reanalysis)	1
sea_surface_temperature_depth	Sea surface temperature adjusted to standard time and depth	1
sst_depth_dtime	Time differences from the base time in the time coordinate variable for the SSTs adjusted to standard depth and time.	1
large_scale_correlated_uncertainty	Systematic uncertainty that is highly correlated between pixels over large scales	1
synoptically_correlated_uncertainty	Systematic uncertainty that is highly correlated between pixels over synoptic scales only	1
uncorrelated_uncertainty	Non-systematic uncertainty (uncorrelated or weakly correlated between pixels)	1
adjustment_uncertainty	Uncertainty in adjustment to standard depth and time (correlated between pixels over synoptic scales)	1
sst_depth_total_uncertainty	To contain the total of the large scale correlated, synoptically correlated, uncorrelated and adjustment uncertainties.	1

4.4.3 Detailed contents

The detailed contents of the products are defined in Table 18. The table specifies the structure of the NetCDF files. Much of this is done with reference to Table 14. However, note that in keeping with the examples in [AD.1] the spatial dimensions are given different names in level 3 files compared to L2P (lat and lon in the former, ni and nj in the latter). Example of how these contents would look in a real file are given in Appendix A.

Table 18. Description of L3 file contents; this follows the specification defined in [AD.1].

Element name	Description	Range/value	Unit	T	D	Bytes
dataset	Begin of dataset					
dimensions	Begin of dimensions					
lon	Zonal dimension.	7200		sl		4
lat	Meridional dimension.	3600		sl		4
time	Time dimension; in practice will be 1.	Unlimited		sl		4
bnds	Dimension for the latitude, longitude and time bounds.	2		sl		4
dimensions	End of dimensions					
variables	Begin of variables					
variable	Begin of variable					
lat	Latitude coordinates	[-90, 90]		fl	lat	lat x 4
attributes	Begin of attributes					
units	Text description of the units.	degrees_north		st		
long_name	A free-text descriptive variable name.	Latitude		st		
standard_name	Unique descriptive name for data.	latitude		st		
valid_min	Minimum valid value for this variable.	-90.0		fl		4
valid_max	Maximum valid value for this variable.	90.0		fl		4
reference_datum	Information about the coordinates.	geographical coordinates, WGS84 projection		st		
axis	Set to X for longitudinal axis, Y for latitudinal axis, T for time.	Y		st		
bounds	Indicates the variable containing the latitude cell boundaries	lat_bnds		st		
comment	Miscellaneous information about the data or the methods used to produce it. (Optional.)			st		
attributes	End of attributes					
variable	End of variable					

Element name	Description	Range/value	Unit	T	D	Bytes
variable	Begin of variable					
lat_bnds	Bounds of the latitude cells	[-90, 90]		fl	bnds x lat	bnds xlat x 4
attributes	Begin of attributes					
units	Text description of the units.	degrees_north		st		
long_name	A free-text descriptive variable name.	Latitude cell boundaries		st		
valid_min	Minimum valid value for this variable.	-90.0		fl		4
valid_max	Maximum valid value for this variable.	90.0		fl		4
comment	Miscellaneous information about the data or the methods used to produce it.	Contains the northern and southern boundaries of the grid cells.		st		
reference_datum	Information about the coordinates.	geographical coordinates, WGS84 projection		st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
lon	Longitude coordinates	[-180, 180]		fl	lon	lon x 4
attributes	Begin of attributes					
units	Text description of the units.	degrees_east		st		
long_name	A free-text descriptive variable name.	Longitude		st		
standard_name	Unique descriptive name for data.	longitude		st		
valid_min	Minimum valid value for this variable.	-180.0		fl		4
valid_max	Maximum valid value for this variable.	180.0		fl		4
reference_datum	Information about the coordinates.	geographical coordinates, WGS84 projection		st		
axis	Set to X for longitudinal axis, Y for latitudinal axis, T for time.	X		st		

Element name	Description	Range/value	Unit	T	D	Bytes
bounds	Indicates the variable containing the longitude cell boundaries	lon_bnds		st		
comment	Miscellaneous information about the data or the methods used to produce it. (Optional.)			st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
lon_bnds	Bounds of the longitude cells	[-180, 180]		fl	bnds x lon	bnds x lon x 4
attributes	Begin of attributes					
units	Text description of the units.	degrees_east		st		
long_name	A free-text descriptive variable name.	Longitude cell boundaries		st		
valid_min	Minimum valid value for this variable.	-180.0		fl		4
valid_max	Maximum valid value for this variable.	180.0		fl		4
comment	Miscellaneous information about the data or the methods used to produce it.	Contains the eastern and western boundaries of the grid cells.		st		
reference_datum	Information about the coordinates.	geographical coordinates, WGS84 projection		st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
time	Time coordinate	[0, 2 ³¹]		sl	time	time x 4
attributes	Begin of attributes					
units	Text description of the units.	seconds since 1981-01-01 00:00:00		st		

Element name	Description	Range/value	Unit	T	D	Bytes
long_name	A free-text descriptive variable name.	reference time of sst file		st		
standard_name	Unique descriptive name for data.	time		st		
axis	Set to X for longitudinal axis, Y for latitudinal axis, T for time.	T		st		
calendar	Defines the calendar used to define the times.	gregorian		st		
bounds	Indicates the variable containing the time cell boundaries	time_bnds		st		
comment	Miscellaneous information about the data or the methods used to produce it. (Optional.)			st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
time_bnds	Bounds of the time cells	[0, 2 ³¹]		sl	bnds x time	bnds x time x 4
attributes	Begin of attributes					
units	Text description of the units.	seconds since 1981-01-01 00:00:00		st		
long_name	A free-text descriptive variable name.	Time cell boundaries		st		
comment	Miscellaneous information about the data or the methods used to produce it.	Contains the start and end times for the time period the data represent.		st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
sea_surface_temperature	As specified in Table 14 (except that the coordinates attribute is optional)					
variable	End of variable					
variable	Begin of variable					

Element name	Description	Range/value	Unit	T	D	Bytes
sst_dtime	As specified in Table 14 (except that the coordinates attribute is optional, the storage type is sl, fill value is -2^{31} and valid min and valid max are -43200 and 43200 respectively since 43200 is half a day in units of seconds)					
variable	End of variable					
variable	Begin of variable					
sses_bias	As specified in Table 14 (except that the coordinates attribute is optional)					
variable	End of variable					
variable	Begin of variable					
sses_standard_deviation	As specified in Table 14 (except that the coordinates attribute is optional)					
variable	End of variable					
variable	Begin of variable					
l2p_flags	As specified in Table 14 (except that the coordinates attribute is optional and if a L3C file do not include 'views' and 'channels' in the flag_meanings attribute and also remove ', 64, 128' from the flag_masks attribute – these are not appropriate for a L3C file since a grid cell could contain data from a mix of retrievals)					
variable	End of variable					
variable	Begin of variable					
quality_level	As specified in Table 14 (except that the coordinates attribute is optional)					
variable	End of variable					
variable	Begin of variable					
wind_speed	As specified in Table 14 (except that the coordinates attribute is optional)					
variable	End of variable					
variable	Begin of variable					
sea_surface_temperature_depth	As specified in Table 14 (except that the coordinates attribute is optional)					
variable	End of variable					
variable	Begin of variable					
sst_depth_dtime	As specified in Table 14 (except that the coordinates attribute is optional, the storage type is sl, fill value is -2^{31} and valid min and valid max are -43200 and 43200 respectively since 43200 is half a day in units of seconds)					
variable	End of variable					
variable	Begin of variable					

Element name	Description	Range/value	Unit	T	D	Bytes
large_scale_correlated_uncertainty	As specified in Table 14 (except that the coordinates attribute is optional)					
variable	End of variable					
variable	Begin of variable					
synoptically_correlated_uncertainty	As specified in Table 14 (except that the coordinates attribute is optional)					
variable	End of variable					
variable	Begin of variable					
uncorrelated_uncertainty	As specified in Table 14 (except that the coordinates attribute is optional)					
variable	End of variable					
variable	Begin of variable					
adjustment_uncertainty	As specified in Table 14 (except that the coordinates attribute is optional)					
variable	End of variable					
variable	Begin of variable					
sst_depth_total_uncertainty	As specified in Table 14 (except that the coordinates attribute is optional)					
variable	End of variable					
variables	End of variables					
attributes	Begin of global attributes					
Global attributes as described in Table 13.						
comment	To contain any miscellaneous information about the data and methods. Delete the 'and sea_surface_temperature_depth' part of the second sentence if the sea_surface_temperature_depth variable is not included in the file.	For information about uncertainty estimates see the comment attributes to the sea_surface_temperature and sea_surface_temperature_depth variables.		st		
northernmost_latitude	Decimal degrees north, range -90 to +90.	90.0		fl		4
geospatial_lat_max	Identical to northernmost_latitude	90.0		fl		4

Element name	Description	Range/value	Unit	T	D	Bytes
southernmost_latitude	Decimal degrees north, range -90 to +90.	-90.0		fl		4
geospatial_lat_min	Identical to southernmost_latitude	-90.0		fl		4
easternmost_longitude	Decimal degrees east, range -180 to +180.	180.0		fl		4
geospatial_lon_max	Identical to easternmost_longitude	180.0		fl		4
westernmost_longitude	Decimal degrees east, range -180 to +180.	-180.0		fl		4
geospatial_lon_min	Identical to westernmost_longitude	-180.0		fl		4
processing_level	Data level.	L3U or L3C		st		
cdm_data_type	"swath" or "grid"	grid		st		
attributes	End of global attributes					
dataset	End of dataset					

4.5 L4 data

4.5.1 Description

Level 4 data – SST data that have been analysed to remove data gaps – are to be produced by the SST_cci project using the Operational Sea surface Temperature and sea Ice Analysis (OSTIA) system. The contents of these files are largely as defined in [AD.1]. A summary of the specification is included here for completeness.

4.5.2 Summary of contents

As described in [AD.1], the contents of the L4 files include SSTs, uncertainties and information about locations of land and sea ice. These are summarised in Table 19.

The contents of the files meets user requirements:

- Provision of a product that does not contain data gaps, such as a L4 product, is required (SST_CCI-UR-QUF-39, [RD.171]).
- SST_CCI-UR-QUF-46 [RD.171]: the product will contain SSTs corresponding to 20 cm depth, which is the joint second most popular choice with potential users.
- SST_CCI-UR-QUF-53 [RD.171]: an estimate of total uncertainty, as provided in the L4 files, is the most popular way of communicating uncertainty with potential users.
- Sea ice fraction and sea ice flags are provided in the L4 files (SST_CCI-UR-QUF-60 and SST_CCI-UR-QUF-69, [RD.171]).

Table 19. Variables to be included within the SST_cci L4 data files; some of the name and description information in this table is reproduced from Section 11.2 of [AD.1]; see [AD.1] for more details. The shading (see Table 7) and the OCC column indicate the number of occurrences of each variable. All variables listed are mandatory (1, shaded blue).

Name	Description	OCC
lat	Central latitude of each spatial point of the data arrays	1
lat_bnds	Latitude bounds of each grid cell; included from [RD.166].	1
lon	Central longitude of each spatial point of the data arrays	1
lon_bnds	Longitude bounds of each grid cell; included from [RD.166].	1
time	Time of each temporal point of the data arrays; time should be the middle of the day that the data represents	1
time_bnds	Time bounds of each time point (start and end of the day); included from [RD.166]	1
analysed_sst	Sea surface temperature data; this will either be a foundation SST (demonstration ECV) or 20 cm depth SST (long term ECV); fill values to be used where there is land; where there is total sea ice cover the analysis will contain temperatures that relax over time towards -1.8°C.	1
analysis_error	Analysis uncertainty (one error standard deviation)	1
sea_ice_fraction	Sea ice concentration; taken from OSI SAF data	1
sea_ice_fraction_error	Estimated error standard deviation of sea ice fraction; taken from OSI SAF data	0..1
mask	Indicates if a location is land, sea-ice or lake	1

4.5.3 Detailed contents

The detailed contents of the files are specified in Table 20. An example of how these contents would look in a real file is given in Appendix A.

Table 20. Detailed contents of the SST_cci L4 files.

Element name	Description	Range/value	Unit	T	D	Bytes
dataset	Begin of dataset					
dimensions	Begin of dimensions					
lon	Zonal dimension	7200		sl	1	4
lat	Meridional dimension	3600		sl	1	4
time	Time dimension; in practice will be 1.	Unlimited		sl	1	4
bnds	Dimension for the latitude, longitude and time bounds.	2		sl		4
dimensions	End of dimensions					
variables	Begin of variables					
variable	Begin of variable					
lat	As Table 18.					
variable	End of variable					
variable	Begin of variable					
lat_bnds	As Table 18.					
variable	End of variable					
variable	Begin of variable					
lon	As Table 18.					
variable	End of variable					
variable	Begin of variable					
lon_bnds	As Table 18.					
variable	End of variable					
variable	Begin of variable					
time	As Table 18.					
variable	End of variable					
variable	Begin of variable					
time_bnds	As Table 18.					
variable	End of variable					
variable	Begin of variable					
analysed_sst	SST values	[270.15, 318.15] (after scaling)	K	ss	time x lat x lon	time x lat x lon x 2

Element name	Description	Range/value	Unit	T	D	Bytes
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-32768		ss		2
units	Text description of the units.	kelvin		st		
scale_factor	To be multiplied by the variable to recover the original value.	0.01		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	273.15		fl		4
long_name	A free-text descriptive variable name.	analysed sea surface temperature		st		
valid_min	Minimum valid value for this variable once they are packed (in storage type).	-300		ss		2
valid_max	Maximum valid value for this variable once they are packed (in storage type).	4500		ss		2
standard_name	If demonstration ECV set to "sea_surface_foundation_temperature" and include the type attribute; if long term ECV set to "sea_water_temperature" and include the depth attribute.	sea_surface_foundation_temperature or sea_water_temperature		st		
source	Set to comma separated list of the values of the global source attribute in the input files; see Section 3.5.			st		
depth	Include only if standard_name is set to sea_water_temperature. Effective depth of the SST data.	20 cm		st		

Element name	Description	Range/value	Unit	T	D	Bytes
comment	Miscellaneous information about the data or the methods used to produce it. (Optional.)			st		
attributes	End of attributes					
variable	End of variable					
variable	Start of variable					
analysis_error	Uncertainty in the SSTs	[0, 327.68] (after scaling)	K	ss	time x lat x lon	time x lat x lon x 2
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-32768		ss		2
units	Text description of the units.	kelvin		st		
scale_factor	To be multiplied by the variable to recover the original value.	0.01		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	0.0		fl		4
long_name	A free-text descriptive variable name.	estimated error standard deviation of analysed_sst		st		
valid_min	Minimum valid value for this variable once they are packed (in storage type).	0		ss		2
valid_max	Maximum valid value for this variable once they are packed (in storage type).	32767		ss		2
standard_name	Unique descriptive name for data.	sea_surface_foundation_temperature_standard_error or sea_surface_water_temperature_standard_error		st		

Element name	Description	Range/value	Unit	T	D	Bytes
comment	Miscellaneous information about the data or the methods used to produce it. (Optional.)			st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
sea_ice_fraction	Sea ice concentration	[0, 1]	Fraction	sc	time x lat x lon	time x lat x lon
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-128		sc		1
units	Text description of the units.	1		st		
scale_factor	To be multiplied by the variable to recover the original value.	0.01		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	0.0		fl		4
long_name	A free-text descriptive variable name.	sea ice area fraction		st		
valid_min	Minimum valid value for this variable once they are packed (in storage type).	0		sc		1
valid_max	Maximum valid value for this variable once they are packed (in storage type).	100		sc		1
standard_name	Unique descriptive name for data.	sea_ice_area_fraction		st		
source	Source of the sea ice concentrations, including version number (to be confirmed).	EUMETSAT_OSI-SAF-ICE-v1.1		st		

Element name	Description	Range/value	Unit	T	D	Bytes
comment	Miscellaneous information about the data or the methods used to produce it. (Optional.)			st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
sea_ice_fraction_error	Sea ice concentration	[0, 1]	Fract ion	sc	time x lat x lon	time x lat x lon
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-128		sc		1
units	Text description of the units.	1		st		
scale_factor	To be multiplied by the variable to recover the original value.	0.01		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	0.0		fl		4
long_name	A free-text descriptive variable name.	sea ice area fraction error estimate		st		
valid_min	Minimum valid value for this variable once they are packed (in storage type).	0		sc		1
valid_max	Maximum valid value for this variable once they are packed (in storage type).	100		sc		1
standard_name	Unique descriptive name for data.	sea_ice_area_fraction_standard_error		st		
source	Source of the sea ice concentrations, including version number (to be confirmed).	EUMETSAT_OSI-SAF-ICE-v1.1		st		

Element name	Description	Range/value	Unit	T	D	Bytes
comment	Miscellaneous information about the data or the methods used to produce it.	Estimated error standard deviation of sea ice fraction		st		
attributes	End of attributes					
variable	End of variable					
variable	Start of variable					
mask	Specifies if a location is land, sea-ice or lake	[1, 31]	None	sc	time x lat x lon	time x lat x lon
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-128		sc		1
long_name	A free-text descriptive variable name.	sea/land/lake/ice field composite mask		st		
valid_min	Minimum valid value for this variable once they are packed (in storage type).	1		sc		1
valid_max	Maximum valid value for this variable once they are packed (in storage type).	31		sc		1
source	Source of the mask, including version numbers.	NAVOCEAN O_landmask_v1.0, EUMETSAT_OSI-SAF_ice mask, ARCLake_lake mask		st		

Element name	Description	Range/value	Unit	T	D	Bytes
comment	Miscellaneous information about the data or methods used to produce them.	b0: 1=grid cell is open sea water b1: 1=grid cell is land b2: 1=grid cell is lake surface b3: 1=grid cell is sea ice b4-b7: reserved for future grid mask data		st		
flag_meanings	Meaning attached to each flag value.	water land optional_lake _surface sea_ice optional_river _surface		st		
flag_masks	Values that correspond to the flags described in flag_meanings.	1b, 2b, 4b, 8b, 16b		sc		1
attributes	End of attributes					
variable	End of variable					
variables	End of variables					
attributes	Begin of global attributes					
As Table 13.						
comment	To contain any miscellaneous information about the data and methods.			st		
northernmost_latitude	Decimal degrees north, range -90 to +90.	90.0		fl		4
geospatial_lat_max	Identical to northernmost_latitude	90.0		fl		4
southernmost_latitude	Decimal degrees north, range -90 to +90.	-90.0		fl		4
geospatial_lat_min	Identical to southernmost_latitude	-90.0		fl		4

Element name	Description	Range/value	Unit	T	D	Bytes
easternmost_longitude	Decimal degrees east, range -180 to +180.	180.0		fl		4
geospatial_lon_max	Identical to easternmost_longitude	180.0		fl		4
westernmost_longitude	Decimal degrees east, range -180 to +180.	-180.0		fl		4
geospatial_lon_min	Identical to westernmost_longitude	-180.0		fl		4
processing_level	Data level.	L4		st		
cdm_data_type	“swath” or “grid”	grid		st		
attributes	End of global attributes					
dataset	End of dataset					

5. FILE SIZE ESTIMATES

An estimate of product size per file for each file type is provided in

Table 21. The product sizes are broken down into components with different dimensionality and the number of bytes used for each component is recorded. For example the table reveals that a level 2 file stores data with dimensionality $n_i \times n_j$ and the total amount of storage required per element of the arrays with these dimensions is 8 bytes (these are the arrays containing the latitudes and longitudes of the data, which are both stored as floating point numbers and use 4 bytes per value). The total storage requirement for these data are therefore $n_i \times n_j \times 8$ bytes, The total file size can be estimated by assuming values for the dimensions and totalling the storage requirements for all the components.

The size estimates give an indication of the data processing and writing requirements. However, they exclude storage for arrays of characters, which use negligible space compared to the data arrays.

Note that these sizes are for uncompressed files. The sizes for L2 data will vary according to sensor but are representative of the largest files that might be expected. In practice the compression capability of the NetCDF-4/HDF5 format will result in much smaller files. However, even if there were no compression the maximum file size would be ~500 MB, which is usable by a high percentage (71%) of users [RD.189].

Table 21. File size estimates in bytes unless otherwise stated, divided into components with different dimensionality. Total size estimate assumes that $n_i = 625$, $n_j = 7,000$, $lon = 7,200$, $lat = 3,600$, $time = 1$, $bnds = 2$. Size of level 2 files will vary from sensor to sensor and level 3/4 file sizes depend on the grid that is used.

Dimensionality	Level 2	Level 3	Level 4
Scalars (attributes)	237	263	176
n_i or lon	0	4	4
n_j or lat	0	4	4
time	4	4	4
$(n_i$ or lon) \times bnds	0	4	4
$(n_j$ or lat) \times bnds	0	4	4
time \times bnds	0	4	4
$(n_i$ or lon) \times (n_j or lat)	8	0	0
$(n_i$ or lon) \times (n_j or lat) \times time	24	28	7
Total uncompressed size estimate	140 MB	726 MB	181 MB

6. ISSUES FOR FUTURE RELEASES OF THIS DOCUMENT

It is recommended that future releases of this product specification should resolve the following issues:

- The institution metadata is specified to be 'ESACCI', which is the RDAC code for the project registered with GRHSST. This is not currently included in the ESA SST CCI Data Standards Working Group vocabulary.

APPENDIX A: CDL LISTINGS OF NETCDF HEADERS

This appendix contains Network Common Data Form Language (CDL) examples of the contents of L2P, L3U and L4 files that conform to the specifications defined in this document.

A.1 L2P example

```
netcdf 20100701003800-ESACCI-L2P_GHRSSST-SSTskin-AVHRRMTA_G-v02.0-
fv01.0.nc {
dimensions:
    ni = 409 ;
    nj = 12348 ;
    time = 1 ;
variables:
    float lat(nj, ni) ;
        lat:units = "degrees_north" ;
        lat:long_name = "Latitude coordinates" ;
        lat:standard_name = "latitude" ;
        lat:valid_min = -90.f ;
        lat:valid_max = 90.f ;
        lat:reference_datum = "geographical coordinates, WGS84
projection" ;
    float lon(nj, ni) ;
        lon:units = "degrees_east" ;
        lon:long_name = "Longitude coordinates" ;
        lon:standard_name = "longitude" ;
        lon:valid_min = -180.f ;
        lon:valid_max = 180.f ;
        lon:reference_datum = "geographical coordinates, WGS84
projection" ;
    int time(time) ;
        time:units = "seconds since 1981-01-01 00:00:00" ;
        time:long_name = "reference time of sst file" ;
        time:standard_name = "time" ;
        time:calendar = "gregorian" ;
    short sea_surface_temperature(time, nj, ni) ;
        sea_surface_temperature:FillValue = -32768s ;
        sea_surface_temperature:units = "kelvin" ;
        sea_surface_temperature:scale_factor = 0.01f ;
        sea_surface_temperature:add_offset = 273.15f ;
        sea_surface_temperature:long_name = "sea surface skin
temperature" ;
        sea_surface_temperature:valid_min = -200s ;
        sea_surface_temperature:valid_max = 5000s ;
        sea_surface_temperature:standard_name =
"sea_surface_skin_temperature" ;
        sea_surface_temperature:comment = "Temperature of the skin
of the ocean; total uncertainty =
sqrt(large_scale_correlated_uncertainty^2+synoptically_correlated_uncert
ainty^2+uncorrelated_uncertainty^2)" ;
        sea_surface_temperature:source = "AVHRRMTA_G-NOAA-L1-v2.0" ;
        sea_surface_temperature:references = "Insert published or
web-based references that describe the data or methods used to produce
them" ;
        sea_surface_temperature:depth = "10 micrometres" ;
        sea_surface_temperature:coordinates = "lon lat" ;
```

```

short sst_dtime(time, nj, ni) ;
sst_dtime:_FillValue = -32768s ;
sst_dtime:units = "seconds" ;
sst_dtime:scale_factor = 1.f ;
sst_dtime:add_offset = 0.f ;
sst_dtime:long_name = "time difference from reference time"
;

sst_dtime:valid_min = -32767s ;
sst_dtime:valid_max = 32767s ;
sst_dtime:comment = "time plus sst_dtime gives seconds after
1981-01-01 00:00:00" ;
sst_dtime:coordinates = "lon lat" ;
byte sses_bias(time, nj, ni) ;
sses_bias:_FillValue = -128b ;
sses_bias:units = "kelvin" ;
sses_bias:scale_factor = 0.01f ;
sses_bias:add_offset = 0.f ;
sses_bias:long_name = "SSES bias estimate" ;
sses_bias:valid_min = -127b ;
sses_bias:valid_max = 127b ;
sses_bias:comment = "Populated with zeroes" ;
sses_bias:coordinates = "lon lat" ;
byte sses_standard_deviation(time, nj, ni) ;
sses_standard_deviation:_FillValue = -128b ;
sses_standard_deviation:units = "kelvin" ;
sses_standard_deviation:scale_factor = 0.01f ;
sses_standard_deviation:add_offset = 1.27f ;
sses_standard_deviation:long_name = "SSES standard
deviation" ;
sses_standard_deviation:valid_min = -127b ;
sses_standard_deviation:valid_max = 127b ;
sses_standard_deviation:comment = "Total uncertainty in each
sea_surface_temperature data point" ;
sses_standard_deviation:coordinates = "lon lat" ;
short l2p_flags(time, nj, ni) ;
l2p_flags:long_name = "L2P flags" ;
l2p_flags:comment = "These flags are important to properly
use the data" ;
l2p_flags:coordinates = "lon lat" ;
l2p_flags:flag_meanings = "microwave land ice lake river
spare views channels" ;
l2p_flags:flag_masks = 1s, 2s, 4s, 8s, 16s, 32s, 64s, 128s ;
byte quality_level(time, nj, ni) ;
quality_level:long_name = "quality level of SST pixel" ;
quality_level:comment = "These are overall quality
indicators and are those used for all GHRSSST SSTs" ;
quality_level:coordinates = "lon lat" ;
quality_level:flag_meanings = "no_data bad_data
worst_quality low_quality acceptable_quality best_quality" ;
quality_level:flag_values = 0b, 1b, 2b, 3b, 4b, 5b ;
byte wind_speed(time, nj, ni) ;
wind_speed:_FillValue = -128b ;
wind_speed:units = "m s-1" ;
wind_speed:scale_factor = 0.1f ;
wind_speed:add_offset = 12.7f ;
wind_speed:long_name = "10m wind speed" ;
wind_speed:valid_min = -127b ;
wind_speed:valid_max = 127b ;
wind_speed:standard_name = "wind_speed" ;

```



```

        wind_speed:comment = "Wind speeds sourced from ECMWF ERA
Interim reanalysis; wind speeds greater than 25.4 m/s are set to 25.4" ;
        wind_speed:source = "ERA_INTERIM-ECMWF-WSP-v1.0" ;
        wind_speed:references = "Insert published or web-based
references that describe the data or methods used to produce them" ;
        wind_speed:coordinates = "lon lat" ;
        wind_speed:height = "10 m" ;
        wind_speed:time_offset = 1.f ;
    short sea_surface_temperature_depth(time, nj, ni) ;
        sea_surface_temperature_depth:_FillValue = -32768s ;
        sea_surface_temperature_depth:units = "kelvin" ;
        sea_surface_temperature_depth:scale_factor = 0.01f ;
        sea_surface_temperature_depth:add_offset = 273.15f ;
        sea_surface_temperature_depth:long_name = "sea surface
temperature at 0.2 m" ;
        sea_surface_temperature_depth:valid_min = -200s ;
        sea_surface_temperature_depth:valid_max = 5000s ;
        sea_surface_temperature_depth:standard_name =
"sea_water_temperature" ;
        sea_surface_temperature_depth:comment = "Temperature of the
ocean at 20 cm depth; total uncertainty =
sqrt(large_scale_correlated_uncertainty^2+synoptically_correlated_uncert
ainty^2+uncorrelated_uncertainty^2+adjustment_uncertainty^2)" ;
        sea_surface_temperature_depth:source = "AVHRRMTA_G-NOAA-L1-
v2.0" ;
        sea_surface_temperature_depth:references = "Insert published
or web-based references that describe the data or methods used to
produce them" ;
        sea_surface_temperature_depth:depth = "0.2 metre" ;
        sea_surface_temperature_depth:coordinates = "lon lat" ;
    short sst_depth_dtime(time, nj, ni) ;
        sst_depth_dtime:_FillValue = -32768s ;
        sst_depth_dtime:units = "seconds" ;
        sst_depth_dtime:scale_factor = 1.f ;
        sst_depth_dtime:add_offset = 0.f ;
        sst_depth_dtime:long_name = "time difference from reference
time" ;
        sst_depth_dtime:valid_min = -32767s ;
        sst_depth_dtime:valid_max = 32767s ;
        sst_depth_dtime:comment = "time plus sst_depth_dtime gives
seconds after 1981-01-01 00:00:00" ;
        sst_depth_dtime:coordinates = "lon lat" ;
    short large_scale_correlated_uncertainty(time, nj, ni) ;
        large_scale_correlated_uncertainty:_FillValue = -32768s ;
        large_scale_correlated_uncertainty:units = "kelvin" ;
        large_scale_correlated_uncertainty:scale_factor = 0.01f ;
        large_scale_correlated_uncertainty:add_offset = 0.f ;
        large_scale_correlated_uncertainty:long_name = "Uncertainty
from errors likely to be correlated over large scales" ;
        large_scale_correlated_uncertainty:valid_min = 0s ;
        large_scale_correlated_uncertainty:valid_max = 5000s ;
        large_scale_correlated_uncertainty:comment = "Component of
uncertainty that is correlated over large scales; can be combined with
other uncertainty estimates to form a total uncertainty" ;
        large_scale_correlated_uncertainty:references = "Insert
published or web-based references that describe the data or methods used
to produce them" ;
        large_scale_correlated_uncertainty:coordinates = "lon lat" ;
    short synoptically_correlated_uncertainty(time, nj, ni) ;
        synoptically_correlated_uncertainty:_FillValue = -32768s ;

```

```

synoptically_correlated_uncertainty:units = "kelvin" ;
synoptically_correlated_uncertainty:scale_factor = 0.01f ;
synoptically_correlated_uncertainty:add_offset = 0.f ;
synoptically_correlated_uncertainty:long_name = "Uncertainty
from errors likely to be correlated over synoptic scales" ;
synoptically_correlated_uncertainty:valid_min = 0s ;
synoptically_correlated_uncertainty:valid_max = 5000s ;
synoptically_correlated_uncertainty:comment = "Component of
uncertainty that is correlated over synoptic scales; can be combined
with other uncertainty estimates to form a total uncertainty" ;
synoptically_correlated_uncertainty:references = "Insert
published or web-based references that describe the data or methods used
to produce them" ;
synoptically_correlated_uncertainty:coordinates = "lon lat"
;
synoptically_correlated_uncertainty:correlation_length_scale
= "100 km" ;
synoptically_correlated_uncertainty:correlation_time_scale =
"1 day" ;
short uncorrelated_uncertainty(time, nj, ni) ;
uncorrelated_uncertainty:_FillValue = -32768s ;
uncorrelated_uncertainty:units = "kelvin" ;
uncorrelated_uncertainty:scale_factor = 0.01f ;
uncorrelated_uncertainty:add_offset = 0.f ;
uncorrelated_uncertainty:long_name = "Uncertainty from
errors likely to be uncorrelated between SSTs" ;
uncorrelated_uncertainty:valid_min = 0s ;
uncorrelated_uncertainty:valid_max = 5000s ;
uncorrelated_uncertainty:comment = "Component of uncertainty
that is uncorrelated between SSTs; can be combined with other
uncertainty estimates to form a total uncertainty" ;
uncorrelated_uncertainty:references = "Insert published or
web-based references that describe the data or methods used to produce
them" ;
uncorrelated_uncertainty:coordinates = "lon lat" ;
short adjustment_uncertainty(time, nj, ni) ;
adjustment_uncertainty:_FillValue = -32768s ;
adjustment_uncertainty:units = "kelvin" ;
adjustment_uncertainty:scale_factor = 0.01f ;
adjustment_uncertainty:add_offset = 0.f ;
adjustment_uncertainty:long_name = "Time and depth
adjustment uncertainty" ;
adjustment_uncertainty:valid_min = 0s ;
adjustment_uncertainty:valid_max = 5000s ;
adjustment_uncertainty:comment = "Adjustment uncertainty;
can be combined with other uncertainty estimates to form a total
uncertainty" ;
adjustment_uncertainty:references = "Insert published or
web-based references that describe the data or methods used to produce
them" ;
adjustment_uncertainty:coordinates = "lon lat" ;
adjustment_uncertainty:correlation_length_scale = "100 km" ;
adjustment_uncertainty:correlation_time_scale = "1 day" ;
short sst_depth_total_uncertainty(time, nj, ni) ;
sst_depth_total_uncertainty:_FillValue = -32768s ;
sst_depth_total_uncertainty:units = "kelvin" ;
sst_depth_total_uncertainty:scale_factor = 0.01f ;
sst_depth_total_uncertainty:add_offset = 0.f ;
sst_depth_total_uncertainty:long_name = "Total uncertainty
in sea_surface_temperature_depth" ;

```

```

sst_depth_total_uncertainty:valid_min = 0s ;
sst_depth_total_uncertainty:valid_max = 5000s ;
sst_depth_total_uncertainty:comment = "Total uncertainty in
each sea_surface_temperature_depth data point" ;
sst_depth_total_uncertainty:coordinates = "lon lat" ;

// global attributes:
:Conventions = "CF-1.5, Unidata Observation Dataset v1.0" ;
:title = "ESA SST CCI AVHRRMTA_G L2P product" ;
:summary = "AVHRRMTA_G L2P product from the ESA SST CCI
project, produced using OE2." ;
:references = "Insert published or web-based references that
describe the data or methods used to produce them" ;
:institution = "ESACCI" ;
:history = "SST CCI processor XXX.YY" ;
:licence = "TBC" ;
:id = "AATSR-ESACCI-L2P-v<Version Number>" ;
:naming_authority = "org.ghrsst" ;
:product_version = "1.0" ;
:uuid = "D7A88FA8-7421-4039-807C-B551D638EDC6" ;
:tracking_id = "D7A88FA8-7421-4039-807C-B551D638EDC6" ;
:gds_version_id = "2.0" ;
:netcdf_version_id = "4.1.1" ;
:date_created = "20110616T152207Z" ;
:file_quality_level = 3 ;
:spatial_resolution = "4 km at nadir" ;
:start_time = "20100701T000000Z" ;
:time_coverage_start = "20100701T000000Z" ;
:stop_time = "20100701T013000Z" ;
:time_coverage_end = "20100701T013000Z" ;
:time_coverage_duration = "PT1H30M" ;
:time_coverage_resolution = "PT1S" ;
:source = "AVHRRMTA_G-NOAA-L1-v2.0, ERA_INTERIM-ECMWF-WSP-
v1.0" ;
:platform = "Envisat" ;
:sensor = "AATSR" ;
:metadata_conventions = "Unidata Dataset Discovery v1.0" ;
:metadata_link = "http://www.esa-cci.org" ;
:keywords = "Oceans > Ocean Temperature > Sea Surface
Temperature" ;
:keywords_vocabulary = "NASA Global Change Master Directory
(GCMD) Science Keywords" ;
:standard_name_vocabulary = "NetCDF Climate and Forecast
(CF) Metadata Convention" ;
:geospatial_lat_units = "degrees_north" ;
:geospatial_lat_resolution = "0.009" ;
:geospatial_lon_units = "degrees_east" ;
:geospatial_lon_resolution = "0.009" ;
:geospatial_vertical_min = -0.2f ;
:geospatial_vertical_max = -1.e-05f ;
:acknowledgment = "Funded by ESA" ;
:creator_name = "ESA SST CCI" ;
:creator_email = "science.leader@esa-sst-cci.org" ;
:creator_url = "http://www.esa-sst-cci.org" ;
:creator_processing_institution = "These data were produced
at <institution> as part of the ESA SST CCI project."
:project = "European Space Agency Sea Surface Temperature
Climate Change Initiative" ;
:publisher_name = "ESACCI" ;

```

```

        :publisher_url = "http://www.esa-sst-cci.org" ;
        :publisher_email = "science.leader@esa-SST-cci.org" ;
        :comment = "See the comment attributes to the
sea_surface_temperature and sea_surface_temperature_depth variables for
information about uncertainty estimates." ;
        :northernmost_latitude = 90.f ;
        :southernmost_latitude = -90.f ;
        :easternmost_longitude = -180.f ;
        :westernmost_longitude = 180.f ;
        :geospatial_lat_max = 90.f ;
        :geospatial_lat_min = -90.f ;
        :geospatial_lon_max = -180.f ;
        :geospatial_lon_min = 180.f ;
        :processing_level = "L2P" ;
        :cdm_data_type = "swath" ;
        :product_specification_version = "SST_CCI-PSD-UKMO-201-
Issue-1-signed" ;
    }

```

A.2 L3U example

```

netcdf 20100701000000-ESACCI-L3U_GHRSSST-SSTskin-AATSR-v02.0-fv01.0 {
dimensions:
    lon = 7200 ;
    lat = 3600 ;
    bnds = 2 ;
    time = UNLIMITED ; // (1 currently)
variables:
    float lat(lat) ;
        lat:units = "degrees_north" ;
        lat:long_name = "Latitude" ;
        lat:standard_name = "latitude" ;
        lat:valid_min = -90.f ;
        lat:valid_max = 90.f ;
        lat:reference_datum = "geographical coordinates, WGS84
projection" ;
        lat:axis = "Y" ;
        lat:bounds = "lat_bnds" ;
    float lat_bnds(lat, bnds) ;
        lat_bnds:units = "degrees_north" ;
        lat_bnds:long_name = "Latitude cell boundaries" ;
        lat_bnds:valid_min = -90.f ;
        lat_bnds:valid_max = 90.f ;
        lat_bnds:comment = "Contains the northern and southern
boundaries of the grid cells." ;
        lat_bnds:reference_datum = "geographical coordinates, WGS84
projection" ;
    float lon(lon) ;
        lon:units = "degrees_east" ;
        lon:long_name = "Longitude" ;
        lon:standard_name = "longitude" ;
        lon:valid_min = -180.f ;
        lon:valid_max = 180.f ;
        lon:reference_datum = "geographical coordinates, WGS84
projection" ;
        lon:axis = "X" ;
        lon:bounds = "lon_bnds" ;
    float lon_bnds(lon, bnds) ;

```

```

lon_bnds:units = "degrees_east" ;
lon_bnds:long_name = "Longitude cell boundaries" ;
lon_bnds:valid_min = -180.f ;
lon_bnds:valid_max = 180.f ;
lon_bnds:comment = "Contains the eastern and western
boundaries of the grid cells." ;
lon_bnds:reference_datum = "geographical coordinates, WGS84
projection" ;
int time(time) ;
time:units = "seconds since 1981-01-01 00:00:00" ;
time:long_name = "reference time of sst file" ;
time:standard_name = "time" ;
time:axis = "T" ;
time:calendar = "gregorian" ;
time:bounds = "time_bnds" ;
int time_bnds(time, bnds) ;
time_bnds:units = "seconds since 1981-01-01 00:00:00" ;
time_bnds:long_name = "Time cell boundaries" ;
time_bnds:comment = "Contains the start and end times for
the time period the data represent" ;
short sea_surface_temperature(time, lat, lon) ;
sea_surface_temperature:_FillValue = -32768s ;
sea_surface_temperature:units = "kelvin" ;
sea_surface_temperature:scale_factor = 0.01f ;
sea_surface_temperature:add_offset = 273.15f ;
sea_surface_temperature:long_name = "sea surface skin
temperature" ;
sea_surface_temperature:valid_min = -200s ;
sea_surface_temperature:valid_max = 5000s ;
sea_surface_temperature:standard_name =
"sea_surface_skin_temperature" ;
sea_surface_temperature:comment = "Temperature of the skin
of the ocean; total uncertainty =
sqrt(large_scale_correlated_uncertainty^2+synoptically_correlated_uncert
ainty^2+uncorrelated_uncertainty^2)" ;
sea_surface_temperature:source = "AATSR-ESA-L1-v2.0" ;
sea_surface_temperature:references = "Insert published or
web-based references that describe the data or methods used to produce
them" ;
sea_surface_temperature:depth = "10 micrometres" ;
int sst_dtime(time, lat, lon) ;
sst_dtime:_FillValue = 0 ;
sst_dtime:units = "seconds" ;
sst_dtime:scale_factor = 1. ;
sst_dtime:add_offset = 0. ;
sst_dtime:long_name = "time difference from reference time"
;
sst_dtime:valid_min = -43200 ;
sst_dtime:valid_max = 43200 ;
sst_dtime:comment = "time plus sst_dtime gives seconds after
1981-01-01 00:00:00" ;
byte sses_bias(time, lat, lon) ;
sses_bias:_FillValue = -128b ;
sses_bias:units = "kelvin" ;
sses_bias:scale_factor = 0.01f ;
sses_bias:add_offset = 1.27f ;
sses_bias:long_name = "SSES bias estimate" ;
sses_bias:valid_min = -127b ;
sses_bias:valid_max = 127b ;
sses_bias:comment = "Populated with zeroes" ;

```

```

byte sses_standard_deviation(time, lat, lon) ;
  sses_standard_deviation:_FillValue = -128b ;
  sses_standard_deviation:units = "kelvin" ;
  sses_standard_deviation:scale_factor = 0.01f ;
  sses_standard_deviation:add_offset = 0.f ;
  sses_standard_deviation:long_name = "SSES standard
deviation" ;
  sses_standard_deviation:valid_min = -127b ;
  sses_standard_deviation:valid_max = 127b ;
  sses_standard_deviation:comment = " Total uncertainty in
each sea_surface_temperature data point" ;
  sses_standard_deviation:coordinates = "lon lat" ;
short l2p_flags(time, lat, lon) ;
  l2p_flags:long_name = "L2P flags" ;
  l2p_flags:comment = "These flags are important to properly
use the data" ;
  l2p_flags:flag_meanings = "microwave land ice lake river
spare views channels" ;
  l2p_flags:flag_masks = 1s, 2s, 4s, 8s, 16s, 32s, 64s, 128s ;
byte quality_level(time, lat, lon) ;
  quality_level:long_name = "quality level of SST pixel" ;
  quality_level:comment = "These are overall quality
indicators and are those used for all GHRSSST SSTs" ;
  quality_level:flag_meanings = "no_data bad_data
worst_quality low_quality acceptable_quality best_quality" ;
  quality_level:flag_values = 0b, 1b, 2b, 3b, 4b, 5b ;
byte wind_speed(time, lat, lon) ;
  wind_speed:_FillValue = -128b ;
  wind_speed:units = "m s-1" ;
  wind_speed:scale_factor = 0.1f ;
  wind_speed:add_offset = 12.7f ;
  wind_speed:long_name = "10m wind speed" ;
  wind_speed:valid_min = -127b ;
  wind_speed:valid_max = 127b ;
  wind_speed:standard_name = "wind_speed" ;
  wind_speed:comment = "Wind speeds sourced from ECMWF ERA
Interim reanalysis; wind speeds greater than 25.4 m/s are set to 25.4" ;
  wind_speed:source = "ERA_INTERIM-ECMWF-WSP-v1.0" ;
  wind_speed:references = "Insert published or web-based
references that describe the data or methods used to produce them" ;
  wind_speed:height = "10 m" ;
  wind_speed:time_offset = 1.f ;
short sea_surface_temperature_depth(time, lat, lon) ;
  sea_surface_temperature_depth:_FillValue = -32768s ;
  sea_surface_temperature_depth:units = "kelvin" ;
  sea_surface_temperature_depth:scale_factor = 0.01f ;
  sea_surface_temperature_depth:add_offset = 273.15f ;
  sea_surface_temperature_depth:long_name = "sea surface
temperature at 0.2 m" ;
  sea_surface_temperature_depth:valid_min = -200s ;
  sea_surface_temperature_depth:valid_max = 5000s ;
  sea_surface_temperature_depth:standard_name =
"sea_water_temperature" ;
  sea_surface_temperature_depth:comment = "Temperature of the
ocean at 20 cm depth; total uncertainty =
sqrt(large_scale_correlated_uncertainty^2+synoptically_correlated_uncert
ainty^2+uncorrelated_uncertainty^2+adjustment_uncertainty^2)" ;
  sea_surface_temperature_depth:source = "AATSR-ESA-L1-v2.0" ;

```

```

        sea_surface_temperature_depth:references = "Insert published
or web-based references that describe the data or methods used to
produce them" ;
        sea_surface_temperature_depth:depth = "0.2 metre" ;
int sst_depth_dtime(time, lat, lon) ;
    sst_depth_dtime:FillValue = 0 ;
    sst_depth_dtime:units = "seconds" ;
    sst_depth_dtime:scale_factor = 1. ;
    sst_depth_dtime:add_offset = 0. ;
    sst_depth_dtime:long_name = "time difference from reference
time" ;
    sst_depth_dtime:valid_min = -43200 ;
    sst_depth_dtime:valid_max = 43200 ;
    sst_depth_dtime:comment = "time plus sst_depth_dtime gives
seconds after 1981-01-01 00:00:00" ;
    short large_scale_correlated_uncertainty(time, lat, lon) ;
    large_scale_correlated_uncertainty:FillValue = -32768s ;
    large_scale_correlated_uncertainty:units = "kelvin" ;
    large_scale_correlated_uncertainty:scale_factor = 0.01f ;
    large_scale_correlated_uncertainty:add_offset = 0.f ;
    large_scale_correlated_uncertainty:long_name = "Uncertainty
from errors likely to be correlated over large scales" ;
    large_scale_correlated_uncertainty:valid_min = 0s ;
    large_scale_correlated_uncertainty:valid_max = 5000s ;
    large_scale_correlated_uncertainty:comment = "Component of
uncertainty that is correlated over large scales; can be combined with
other uncertainty estimates to form a total uncertainty" ;
    large_scale_correlated_uncertainty:references = "Insert
published or web-based references that describe the data or methods used
to produce them" ;
    short synoptically_correlated_uncertainty(time, lat, lon) ;
    synoptically_correlated_uncertainty:FillValue = -32768s ;
    synoptically_correlated_uncertainty:units = "kelvin" ;
    synoptically_correlated_uncertainty:scale_factor = 0.01f ;
    synoptically_correlated_uncertainty:add_offset = 0.f ;
    synoptically_correlated_uncertainty:long_name = "Uncertainty
from errors likely to be correlated over synoptic scales" ;
    synoptically_correlated_uncertainty:valid_min = 0s ;
    synoptically_correlated_uncertainty:valid_max = 5000s ;
    synoptically_correlated_uncertainty:comment = "Component of
uncertainty that is correlated over synoptic scales; can be combined
with other uncertainty estimates to form a total uncertainty" ;
    synoptically_correlated_uncertainty:references = "Insert
published or web-based references that describe the data or methods used
to produce them" ;
    synoptically_correlated_uncertainty:correlation_length_scale
= "100 km" ;
    synoptically_correlated_uncertainty:correlation_time_scale =
"1 day" ;
    short uncorrelated_uncertainty(time, lat, lon) ;
    uncorrelated_uncertainty:FillValue = -32768s ;
    uncorrelated_uncertainty:units = "kelvin" ;
    uncorrelated_uncertainty:scale_factor = 0.01f ;
    uncorrelated_uncertainty:add_offset = 0.f ;
    uncorrelated_uncertainty:long_name = "Uncertainty from
errors likely to be uncorrelated between SSTs" ;
    uncorrelated_uncertainty:valid_min = 0s ;
    uncorrelated_uncertainty:valid_max = 5000s ;

```

```

        uncorrelated_uncertainty:comment = "Component of uncertainty
that is uncorrelated between SSTs; can be combined with other
uncertainty estimates to form a total uncertainty" ;
        uncorrelated_uncertainty:references = "Insert published or
web-based references that describe the data or methods used to produce
them" ;
        short adjustment_uncertainty(time, lat, lon) ;
        adjustment_uncertainty:_FillValue = -32768s ;
        adjustment_uncertainty:units = "kelvin" ;
        adjustment_uncertainty:scale_factor = 0.01f ;
        adjustment_uncertainty:add_offset = 0.f ;
        adjustment_uncertainty:long_name = "Time and depth
adjustment uncertainty" ;
        adjustment_uncertainty:valid_min = 0s ;
        adjustment_uncertainty:valid_max = 5000s ;
        adjustment_uncertainty:comment = "Adjustment uncertainty;
can be combined with other uncertainty estimates to form a total
uncertainty" ;
        adjustment_uncertainty:references = "Insert published or
web-based references that describe the data or methods used to produce
them" ;
        adjustment_uncertainty:correlation_length_scale = "100 km" ;
        adjustment_uncertainty:correlation_time_scale = "1 day" ;
        short sst_depth_total_uncertainty(time, nj, ni) ;
        sst_depth_total_uncertainty:_FillValue = -32768s ;
        sst_depth_total_uncertainty:units = "kelvin" ;
        sst_depth_total_uncertainty:scale_factor = 0.01f ;
        sst_depth_total_uncertainty:add_offset = 0.f ;
        sst_depth_total_uncertainty:long_name = "Total uncertainty
in sea_surface_temperature_depth" ;
        sst_depth_total_uncertainty:valid_min = 0s ;
        sst_depth_total_uncertainty:valid_max = 5000s ;
        sst_depth_total_uncertainty:comment = "Total uncertainty in
each sea_surface_temperature_depth data point" ;

// global attributes:
:Conventions = "CF-1.5, Unidata Observation Dataset v1.0" ;
:title = "ESA SST CCI AATSR L3U product" ;
:summary = "AATSR L3U product from the ESA SST CCI project,
produced using <algorithm name>." ;
:references = "Insert published or web-based references that
describe the data or methods used to produce them" ;
:institution = "ESACCI" ;
:history = "SST CCI processor XXX.YY" ;
:licence = "TBC" ;
:id = "AATSR-ESACCI-L3U-v<Version Number>" ;
:naming_authority = "org.ghrsst" ;
:product_version = "1.0" ;
:uuid = "D7A88FA8-7421-4039-807C-B551D638EDC6" ;
:tracking_id = "D7A88FA8-7421-4039-807C-B551D638EDC6" ;
:gds_version_id = "2.0" ;
:netcdf_version_id = "4.1.1" ;
:date_created = "20110616T152612Z" ;
:file_quality_level = 3 ;
:spatial_resolution = "1.1km at nadir" ;
:start_time = "20100701T000000Z" ;
:time_coverage_start = "20100701T000000Z" ;
:stop_time = "20100701T013000Z" ;
:time_coverage_end = "20100701T013000Z" ;
:time_coverage_duration = "PT1H30M" ;

```



```

:time_coverage_resolution = "PT1S" ;
:source = "AATSR-ESA-L1-v2.0, ERA_INTERIM-ECMWF-WSP-v1.0" ;
:platform = "Envisat" ;
:sensor = "AATSR" ;
:metadata_conventions = "Unidata Dataset Discovery v1.0" ;
:metadata_link = "http://www.esa-cci.org" ;
:keywords = "Oceans > Ocean Temperature > Sea Surface
Temperature" ;
:keywords_vocabulary = "NASA Global Change Master Directory
(GCMD) Science Keywords" ;
:standard_name_vocabulary = "NetCDF Climate and Forecast
(CF) Metadata Convention" ;
:geospatial_lat_units = "degrees_north" ;
:geospatial_lat_resolution = "0.05" ;
:geospatial_lon_units = "degrees_east" ;
:geospatial_lon_resolution = "0.05" ;
:geospatial_vertical_min = -0.2f ;
:geospatial_vertical_max = -1.e-05f ;
:acknowledgment = "Funded by ESA" ;
:creator_name = "ESA SST CCI" ;
:creator_email = "science.leader@esa-sst-cci.org" ;
:creator_url = "http://www.esa-sst-cci.org" ;
:creator_processing_institution = "These data were produced
at <institution> as part of the ESA SST CCI project."
:project = "European Space Agency Sea Surface Temperature
Climate Change Initiative" ;
:publisher_name = "ESACCI" ;
:publisher_url = "http://www.esa-sst-cci.org" ;
:publisher_email = "science.leader@esa-SST-cci.org" ;
:comment = "See the comment attributes to the
sea_surface_temperature and sea_surface_temperature_depth variables for
information about uncertainty estimates." ;
:northernmost_latitude = 90.f ;
:southernmost_latitude = -90.f ;
:easternmost_longitude = -180.f ;
:westernmost_longitude = 180.f ;
:geospatial_lat_max = 90.f ;
:geospatial_lat_min = -90.f ;
:geospatial_lon_max = -180.f ;
:geospatial_lon_min = 180.f ;
:processing_level = "L3U" ;
:cdm_data_type = "grid" ;
:product_specification_version = "SST_CCI-PSD-UKMO-201-
Issue-1-signed" ;
}

```

A.3 L4 example

```

netcdf 20100701000000-ESACCI-L4_GHRSSST-SSTdepth-OSTIA-GLOB-v02.0-fv01.0
{
dimensions:
    lon = 7200 ;
    lat = 3600 ;
    bnds = 2 ;
    time = UNLIMITED ; // (1 currently)
variables:
    float lat(lat) ;
        lat:units = "degrees_north" ;
        lat:long_name = "Latitude" ;

```

```

        lat:standard_name = "latitude" ;
        lat:valid_min = -90.f ;
        lat:valid_max = 90.f ;
        lat:reference_datum = "geographical coordinates, WGS84
projection" ;
        lat:axis = "Y" ;
        lat:bounds = "lat_bnds" ;
        float lat_bnds(lat, bnds) ;
        lat_bnds:units = "degrees_north" ;
        lat_bnds:long_name = "Latitude cell boundaries" ;
        lat_bnds:valid_min = -90.f ;
        lat_bnds:valid_max = 90.f ;
        lat_bnds:comment = "Contains the northern and southern
boundaries of the grid cells." ;
        lat_bnds:reference_datum = "geographical coordinates, WGS84
projection" ;
        float lon(lon) ;
        lon:units = "degrees_east" ;
        lon:long_name = "Longitude" ;
        lon:standard_name = "longitude" ;
        lon:valid_min = -180.f ;
        lon:valid_max = 180.f ;
        lon:reference_datum = "geographical coordinates, WGS84
projection" ;
        lon:axis = "X" ;
        lon:bounds = "lon_bnds" ;
        float lon_bnds(lon, bnds) ;
        lon_bnds:units = "degrees_east" ;
        lon_bnds:long_name = "Longitude cell boundaries" ;
        lon_bnds:valid_min = -180.f ;
        lon_bnds:valid_max = 180.f ;
        lon_bnds:comment = "Contains the eastern and western
boundaries of the grid cells." ;
        lon_bnds:reference_datum = "geographical coordinates, WGS84
projection" ;
        int time(time) ;
        time:units = "seconds since 1981-01-01 00:00:00" ;
        time:long_name = "reference time of sst file" ;
        time:standard_name = "time" ;
        time:axis = "T" ;
        time:calendar = "gregorian" ;
        time:bounds = "time_bnds" ;
        float time_bnds(time, bnds) ;
        time_bnds:units = "seconds since 1981-01-01 00:00:00" ;
        time_bnds:long_name = "Time cell boundaries" ;
        time_bnds:comment = "Contains the start and end times for
the time period the data represent" ;
        short analysed_sst(time, lat, lon) ;
        analysed_sst:FillValue = -32768s ;
        analysed_sst:units = "kelvin" ;
        analysed_sst:scale_factor = 0.01f ;
        analysed_sst:add_offset = 273.15f ;
        analysed_sst:long_name = "analysed sea surface temperature"
;
        analysed_sst:valid_min = -300s ;
        analysed_sst:valid_max = 4500s ;
        analysed_sst:standard_name = "sea_water_temperature" ;
        analysed_sst:source = "AATSR-ESACCI-L3U-LT-v01.0, AVHRR19_G-
ESACCI-L2P-LT-v01.0" ;
        analysed_sst:depth = "20 cm" ;

```

```

short analysis_error(time, lat, lon) ;
analysis_error:FillValue = -32768s ;
analysis_error:units = "kelvin" ;
analysis_error:scale_factor = 0.01f ;
analysis_error:add_offset = 0.f ;
analysis_error:long_name = "estimated error standard
deviation of analysed_sst" ;
analysis_error:valid_min = 0s ;
analysis_error:valid_max = 32767s ;
analysis_error:standard_name = "sea_water_temperature
standard_error" ;
byte sea_ice_fraction(time, lat, lon) ;
sea_ice_fraction:FillValue = -128b ;
sea_ice_fraction:units = "1" ;
sea_ice_fraction:scale_factor = 0.01f ;
sea_ice_fraction:add_offset = 0.f ;
sea_ice_fraction:long_name = "sea ice area fraction" ;
sea_ice_fraction:valid_min = 0s ;
sea_ice_fraction:valid_max = 100s ;
sea_ice_fraction:standard_name = "sea_ice_area_fraction" ;
sea_ice_fraction:source = "EUMETSAT_OSI-SAF-ICE-v1.1" ;
byte sea_ice_fraction_error(time, lat, lon) ;
sea_ice_fraction:FillValue = -128b ;
sea_ice_fraction:units = "1" ;
sea_ice_fraction:scale_factor = 0.01f ;
sea_ice_fraction:add_offset = 0.f ;
sea_ice_fraction:long_name = "sea ice area fraction error
estimate" ;
sea_ice_fraction:valid_min = 0s ;
sea_ice_fraction:valid_max = 100s ;
sea_ice_fraction:standard_name = "sea_ice_area_fraction
standard_error" ;
sea_ice_fraction:source = "EUMETSAT_OSI-SAF-ICE-v1.1" ;
byte mask(time, lat, lon) ;
mask:FillValue = -128b ;
mask:long_name = "sea/land/lake/ice field composite mask" ;
mask:valid_min = 1b ;
mask:valid_max = 31b ;
mask:source = "NAVOCEANO_landmask_v1.0, EUMETSAT_OSI-
SAF_icemask, ARCLake_lakemask" ;
mask:comment = "b0: 1=grid cell is open sea water b1: 1=grid
cell is land b2: 1=grid cell is lake surface b3: 1=grid cell is sea ice
b4-b7: reserved for future grid mask data" ;
mask:flag_meanings = "water land optional_lake_surface sea-
ice optional_river_surface" ;
mask:flag_masks = 1b, 2b, 4b, 8b, 16b ;

// global attributes:
:Conventions = "CF-1.5, Unidata Observation Dataset v1.0" ;
:title = "ESA SST CCI OSTIA L4 product" ;
:summary = "OSTIA L4 product from the ESA SST CCI project,
produced using <algorithm name>." ;
:references = "Insert published or web-based references that
describe the data or methods used to produce them" ;
:institution = "ESACCI" ;
:history = "SST CCI processor XXX.YY" ;
:licence = "TBC" ;
:id = "OSTIA-ESACCI-L4-v<Version Number>" ;
:naming_authority = "org.ghrsst" ;
:product_version = "1.0" ;

```

```

:uuid = "D7A88FA8-7421-4039-807C-B551D638EDC6" ;
:tracking_id = "D7A88FA8-7421-4039-807C-B551D638EDC6" ;
:gds_version_id = "2.0" ;
:netcdf_version_id = "4.1.1" ;
:date_created = "20110616T152836Z" ;
:file_quality_level = 3 ;
:spatial_resolution = "1.1km at nadir" ;
:start_time = "20100701T000000Z" ;
:time_coverage_start = "20100701T000000Z" ;
:stop_time = "20100701T235959Z" ;
:time_coverage_end = "20100701T235959Z" ;
:time_coverage_duration = "P1D" ;
:time_coverage_resolution = "P1D" ;
:source = "AATSR-ESACCI-L3U-v1.0, AVHRR-ESACCI-L2P-v1.0,
EUMETSAT_OSI-SAF-ICE-v1.1" ;
:platform = "Envisat, NOAA-<X>" ;
:sensor = "AATSR" ;
:metadata_conventions = "Unidata Dataset Discovery v1.0" ;
:metadata_link = "http://www.esa-cci.org" ;
:keywords = "Oceans > Ocean Temperature > Sea Surface
Temperature" ;
:keywords_vocabulary = "NASA Global Change Master Directory
(GCMD) Science Keywords" ;
:standard_name_vocabulary = "NetCDF Climate and Forecast
(CF) Metadata Convention" ;
:geospatial_lat_units = "degrees_north" ;
:geospatial_lat_resolution = "0.05" ;
:geospatial_lon_units = "degrees_east" ;
:geospatial_lon_resolution = "0.05" ;
:geospatial_vertical_min = -0.2f ;
:geospatial_vertical_max = -0.2f ;
:acknowledgment = "Funded by ESA" ;
:creator_name = "ESA SST CCI" ;
:creator_email = "science.leader@esa-sst-cci.org" ;
:creator_url = "http://www.esa-sst-cci.org" ;
:creator_processing_institution = "These data were produced
at the Met Office as part of the ESA SST CCI project."
:project = "European Space Agency Sea Surface Temperature
Climate Change Initiative" ;
:publisher_name = "ESACCI" ;
:publisher_url = "http://www.esa-sst-cci.org" ;
:publisher_email = "science.leader@esa-SST-cci.org" ;
:comment = "" ;
:northernmost_latitude = 90.f ;
:southernmost_latitude = -90.f ;
:easternmost_longitude = -180.f ;
:westernmost_longitude = 180.f ;
:geospatial_lat_max = 90.f ;
:geospatial_lat_min = -90.f ;
:geospatial_lon_max = -180.f ;
:geospatial_lon_min = 180.f ;
:processing_level = "L4" ;
:cdm_data_type = "grid" ;
:product_specification_version = "SST_CCI-PSD-UKMO-201-
Issue-1-signed" ;
}

```

APPENDIX B: SPECIFICATION OF FILES TO STORE SSTs FROM INDIVIDUAL RETRIEVALS

The SST_cci products will contain SSTs derived using the best available retrievals. For example a retrieval that makes use of three channels would be preferred to a two channel retrieval. No user requirement has been identified for the 'non-best' retrievals. However, it is possible that a user requirement may arise and so a specification for files to hold these retrievals is presented here. These are referred to as auxiliary files.

Auxiliary files are defined to contain each individual retrieval type and the adjustments that can be used to translate the SSTs to represent a standard depth and time (produced for the long term ECV only). As defined here, these can only be used in combination with the SST_cci files containing the best retrievals as they do not contain variables that do not vary with retrieval type. By not including these variables, storage requirements are reduced. The user would obtain any data and metadata that are not included in the auxiliary file from the file containing the best retrievals. However, this clearly has the disadvantage that the complexity of managing the data is increased.

B.1 L2P auxiliary files

Two basic auxiliary file structures are specified as containers for:

1. types of retrievals additional to the 'best available' retrievals in the L2P files, and,
2. adjustments to convert the SSTs to a standard depth and time (20 cm depth at 10.30 am or pm).

The first type of auxiliary file will store the results from different (alternative) retrieval algorithms. The adjustments to convert to standard depth and time are independent of retrieval type and so are stored separately in the second type of auxiliary file. The contents of these are designed to be minimal to reduce storage space. They are only usable in combination with their corresponding L2P file. Attributes in the file informs the user of the name of the full L2P file that corresponds to the auxiliary file and its universally unique identifier.

Files for the different retrievals

File name

The file name shall be identical to the name of the L2P file except the <Additional Segregator> component shall additionally be used to distinguish between retrievals. The form of this shall be:

<Additional Segregator> = <LT or DM>_Aux_<Retrieval_><View_><Channels>

<Retrieval_> is used to record the nature of the retrieval algorithm; for example "OE1_", "OE2_".

<View_> is used to record the number of views used in the retrieval; possible values are "Nadir_" and "Dual_". Only include this for ATSR data as it is not applicable to other sensors.

<Channels> is used to record the number of channels used in the retrieval; possible values are "2_channel" and "3_channel".

File structure

The structure of the file is described in Table 22. It comprises components that have changed relative to the L2P file. The adjusted SST for the alternative retrieval can be calculated by the user from the information in the auxiliary file containing the adjustments. Global attributes are provided to link the file to its L2P counterpart and to the file containing the adjustments using with the filenames or the universally unique identifiers. The full set of global attributes from the full L2P file is also included.

Table 22. Specification of auxiliary files that can be used to store alternative retrievals to those held in the L2P file.

Element name	Description	Range/value	Unit	T	D	Bytes
dataset	Begin of dataset					
dimensions	Begin of dimensions					
ni	Across track dimension.			sl	1	4
nj	Along track dimension.			sl	1	4
time	Time dimension; must be 1 for L2P data.	1		sl	1	4
dimensions	End of dimensions					
variables	Begin of variables					
variable	Begin of variable					
sea_surface_temperature	Sea surface temperature retrievals	[271.15, 323.15] (after scaling)	K	ss	time, nj, ni	time × nj × ni × 2
attributes	Beginning of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-32768		ss		2
units	Text description of the units.	kelvin		st		
scale_factor	To be multiplied by the variable to recover the original value.	0.01		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	273.15		fl		4
long_name	A free-text descriptive variable name. Insert	sea surface skin temperature				

Element name	Description	Range/value	Unit	T	D	Bytes
	algorithm name as appropriate.	retrieved using <algorithm>				
standard_name	Unique descriptive name for data.	sea_surface_skin_temperature				
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
l2p_flags	Specifies type of input data and provide flags and information to the user.	[0, 255] (this may change if extra flags are defined)		ss	time x nj x ni	time x nj x ni x 2
attributes	Beginning of attributes					
long_name	A free-text descriptive variable name.	L2P flags		st		
flag_meanings	Meaning attached to each flag value.	microwave land ice lake river spare views channels		st		
flag_masks	Bit masks corresponding to the flags described in flag_meanings.	1, 2, 4, 8, 16, 32, 64, 128		ss		2
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
quality_level	Indicator of the quality of each SST.	[0, 5]		sc	time x nj x ni	time x nj x ni
attributes	Beginning of attributes					
long_name	A free-text descriptive variable name.	quality level of SST pixel		st		
flag_meanings	Meaning attached to each flag value.	no_data bad_data worst_quality low_quality acceptable_quality best_quality		st		
flag_values	Values that	0b, 1b, 2b,		sc		1

Element name	Description	Range/value	Unit	T	D	Bytes
	correspond to the flags described in flag_meanings. The b after the numbers is to indicate that they are single byte numbers.	3b, 4b, 5b				
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
large_scale_correlated_uncertainty	Component of uncertainty that is highly correlated between SST retrievals separated by larger than sub-synoptic spatio-temporal scales.	[0, 50] (after scaling)	K	ss	time x nj x ni	time x nj x ni x 2
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-32768		ss		2
units	Text description of the units.	kelvin		st		
scale_factor	To be multiplied by the variable to recover the original value.	0.01		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	0.0		fl		4
long_name	A free-text descriptive variable name.	Uncertainty from errors likely to be correlated over large scales		st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
synoptically_correlated_uncertainty	Component of uncertainty that is	[0, 50] (after scaling)	K	ss	time x nj	time x nj x ni

Element name	Description	Range/value	Unit	T	D	Bytes
	highly correlated between SST retrievals separated by sub-synoptic spatio-temporal scales.				x ni	x 2
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-32768		ss		2
units	Text description of the units.	kelvin		st		
scale_factor	To be multiplied by the variable to recover the original value.	0.01		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	0.0		fl		4
long_name	A free-text descriptive variable name.	Uncertainty from errors like to be correlated over synoptic scales		st		
correlation_length_scale	Estimate of the correlation length scale of the uncertainties.	100 km		st		
correlation_time_scale	Estimate of the correlation time scale of the uncertainties	1 day		st		
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
uncorrelated_uncertainty	Component of uncertainty that is uncorrelated between SST retrievals separated by any spatio-temporal scales.	[0, 50] (after scaling)	K	ss	time x nj x ni	time x nj x ni x 2

Element name	Description	Range/value	Unit	T	D	Bytes
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-32768		ss		2
units	Text description of the units.	kelvin		st		
scale_factor	To be multiplied by the variable to recover the original value.	0.01		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	0.0		fl		4
long_name	A free-text descriptive variable name.	Uncertainty from errors likely to be uncorrelated between SSTs		st		
attributes	End of attributes					
variable	End of variable					
variables	End of variables					
attributes	Begin of global attribute					
Global attributes as for the full L2P files (see Table 14).						
full_file	The name of the L2P file that these data correspond to.			st		
full_file_uuid	The universally unique identifier of the full file.					
adj_file	The name of the auxiliary file containing the adjustments to standard depth and time and the corresponding uncertainties. Only needs to be inserted where tos is included in the L2P file.			st		

Element name	Description	Range/value	Unit	T	D	Bytes
adj_file_uuid	The universally unique identifier of the adjustments auxiliary file.			st		
attributes	End of global attributes					
dataset	End of dataset					

Files for the adjustments

File name

The file name shall be identical to the name of the L2P file except the <Additional Segregator> component shall additionally be used to distinguish between retrievals. The form of this shall be:

<Additional Segregator> = <LT or DM>_Adj_<Algorithm_>00.2m_10.30

<Algorithm_> is used to for text that distinguishes between different methods of obtaining adjustments.

File structure

The structure of the file is described in Table 23. It comprises only the adjustments to convert the SSTs to the standard depth and time. Adjustments shall be defined anywhere that an SST retrieval is possible. Global attributes are provided to link the file to its L2P counterpart using either the file name or the universally unique identifier. The global attributes from the L2P file are all also included.

Table 23. Specification of auxiliary files that can be used to store adjustments convert SSTs to a standard depth and time.

Element name	Description	Range/value	Unit	T	D	Bytes
dataset	Begin of dataset					
dimensions	Begin of dimensions					
ni	Across track dimension.	7200		sl	1	4
nj	Along track dimension.	3600		sl	1	4
time	Time dimension; must be 1 for L2P data.	1		sl	1	4
dimensions	End of dimensions					
variables	Begin of variables					
variable	Begin of variable					
sst_adjustments		[-32767, 32767]	K	ss	time, nj, ni	time x nj x ni

Element name	Description	Range/value	Unit	T	D	Bytes
						x 2
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-32768		ss		2
units	Text description of the units.	kelvin		st		
scale_factor	To be multiplied by the variable to recover the original value.	0.01		fl		4
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	0.0		fl		4
long_name	A free-text descriptive variable name.	Adjustment for SSTs to make them correspond to 10.30 local time and depth of 20 cm				
attributes	End of attributes					
variable	End of variable					
variable	Begin of variable					
adjustment_uncertainty	Uncertainty in adjustment for depth and time, correlated between SST estimates separated by sub-synoptic spatio-temporal scales.	[0, 50] (after scaling)	K	ss	time x nj x ni	time x nj x ni x 2
attributes	Begin of attributes					
_FillValue	A value used to indicate array elements containing no valid data.	-32768		ss		2
units	Text description of the units.	kelvin		st		
scale_factor	To be multiplied by the variable to recover the original	0.01		fl		4

Element name	Description	Range/value	Unit	T	D	Bytes
	value.					
add_offset	To be added to the variable after multiplying by the scale factor to recover the original value.	0.0		fl		4
long_name	A free-text descriptive variable name.	Time and depth adjustment uncertainty		st		
correlation_length_scale	Estimate of the correlation length scale of the uncertainties.	100 km		st		
correlation_time_scale	Estimate of the correlation time scale of the uncertainties	1 day		st		
attributes	End of attributes					
variable	End of variable					
variables	End of variables					
attributes	Begin of global attribute					
Global attributes as for the full L2P files (see Table 14).						
full_file	The name of the L2P file that these data correspond to.			st		
adj_file_uuid	The universally unique identifier of the adjustments auxiliary file.			st		
attributes	End of global attributes					
dataset	End of dataset					

B.2 L3 auxiliary files

The SST_cci L3 files (L3U and L3C) are to hold the best available retrievals of SST. As with L2 data, auxiliary files are defined to hold other retrievals and the adjustment to standard depth and time. The same definitions as for L2 data are used; these are defined in Table 22 and Table 23. As with the L2P and L3 file specification, dimensions are named lon and lat in place of ni and nj. The global attributes shall follow the L3 definition given in Table 18, rather than the L2P definitions from Table 14.

B.2.1 Auxiliary file size estimates

Estimates of the sizes of the auxiliary files are given in

Table 24. See Section 5 for more details about the contents of this table.

Table 24. File size estimates in bytes unless otherwise stated, divided into components with different dimensionality. Total size estimate assumes that $n_i = 512$, $n_j = 28,000$, $lon = 7,200$, $lat = 3,600$, $time = 1$, $bnds = 2$. Size of level 2 files will vary from sensor to sensor and level 3 file sizes depend on the grid that is used.

Dimensionality	Level 2 auxiliary – alternate retrievals	Level 2 auxiliary – adjustments	Level 3 auxiliary – alternate retrievals	Level 3 auxiliary – adjustments
Scalars (attributes)	107	84	107	84
n_i or lon	0	0	0	0
n_j or lat	0	0	0	0
time	0	0	0	0
$(n_i$ or $lon) \times bnds$	0	0	0	0
$(n_j$ or $lat) \times bnds$	0	0	0	0
time \times bnds	0	0	0	0
$(n_i$ or $lon) \times (n_j$ or $lat)$	0	0	0	0
$(n_i$ or $lon) \times (n_j$ or $lat) \times$ time	11	4	11	4
Total uncompressed size estimate	150 MB	55 MB	272 MB	99 MB

B.2.2 CDL listings of NetCDF headers for the auxiliary files

As described in Appendix A, CDL listings of the NetCDF headers of example files are provided.

B.2.2.1 L2P auxiliary file containing SST retrievals

```
netcdf 20100701000000-ESACCI-L2P_GHRSSST-SSTskin-
AVHRRMTA_G_Aux_OE1_Nadir_2_channel-v02.0-fv01.0 {
dimensions:
    ni = 512 ;
    nj = 28000 ;
    time = 1 ;
variables:
    short sea_surface_temperature(time, nj, ni) ;
    sea_surface_temperature:_FillValue = -32768s ;
    sea_surface_temperature:units = "kelvin" ;
    sea_surface_temperature:scale_factor = 0.01f ;
    sea_surface_temperature:add_offset = 273.15f ;
    sea_surface_temperature:long_name = "sea surface temperature
retrieved using 2-channels, dual view" ;
    sea_surface_temperature:standard_name =
"sea_surface_skin_temperature" ;
    short l2p_flags(time, nj, ni) ;
```

```

        l2p_flags:long_name = "L2P flags" ;
        l2p_flags:flag_meanings = "microwave land ice lake river
spare views channel" ;
        l2p_flags:flag_masks = 1s, 2s, 4s, 8s, 16s, 32s, 64s, 128s ;
byte quality_level(time, nj, ni) ;
    quality_level:long_name = "quality level of SST pixel" ;
    quality_level:flag_meanings = "no_data bad_data
worst_quality low_quality acceptable_quality best_quality" ;
    quality_level:flag_values = 0b, 1b, 2b, 3b, 4b, 5b ;
short large_scale_correlated_uncertainty(time, nj, ni) ;
    large_scale_correlated_uncertainty:_FillValue = -32768s ;
    large_scale_correlated_uncertainty:units = "kelvin" ;
    large_scale_correlated_uncertainty:scale_factor = 0.01f ;
    large_scale_correlated_uncertainty:add_offset = 0.f ;
    large_scale_correlated_uncertainty:long_name = "Uncertainty
from errors likely to be correlated over large scales" ;
short synoptically_correlated_uncertainty(time, nj, ni) ;
    synoptically_correlated_uncertainty:_FillValue = -32768s ;
    synoptically_correlated_uncertainty:units = "kelvin" ;
    synoptically_correlated_uncertainty:scale_factor = 0.01f ;
    synoptically_correlated_uncertainty:add_offset = 0.f ;
    synoptically_correlated_uncertainty:long_name = "Uncertainty
from errors likely to be correlated over synoptic scales" ;
    synoptically_correlated_uncertainty:correlation_length_scale
= "100 km" ;
    synoptically_correlated_uncertainty:correlation_time_scale =
"1 day" ;
short uncorrelated_uncertainty(time, nj, ni) ;
    uncorrelated_uncertainty:_FillValue = -32768s ;
    uncorrelated_uncertainty:units = "kelvin" ;
    uncorrelated_uncertainty:scale_factor = 0.01f ;
    uncorrelated_uncertainty:add_offset = 0.f ;
    uncorrelated_uncertainty:long_name = "Uncertainty from
errors likely to be uncorrelated between SSTs" ;

// global attributes:
    :Conventions = "CF-1.5, Unidata Observation Dataset v1.0" ;
    :title = "ESA SST CCI AATSR L2P product" ;
    :summary = "AVHRRMTA_G L2P product from the ESA SST CCI
project, produced using <algorithm name>." ;
    :references = "Insert published or web-based references that
describe the data or methods used to produce them" ;
    :institution = "ESACCI" ;
    :history = "SST CCI processor XXX.YY" ;
    :licence = "TBC" ;
    :id = "AVHRRMTA_G-ESACCI-L2P-v<Version Number>" ;
    :naming_authority = "org.ghrsst" ;
    :product_version = "1.0" ;
    :uuid = "E7A88FA8-7421-4039-807C-B551D638EDC6" ;
    :tracking_id = "E7A88FA8-7421-4039-807C-B551D638EDC6" ;
    :gds_version_id = "2.0" ;
    :netcdf_version_id = "4.1.1" ;
    :date_created = "20110616T152318Z" ;
    :file_quality_level = 3 ;
    :spatial_resolution = "4 km at nadir" ;
    :start_time = "20100701T000000Z" ;
    :time_coverage_start = "20100701T000000Z" ;
    :stop_time = "20100701T013000Z" ;
    :time_coverage_end = "20100701T013000Z" ;
    :time_coverage_duration = "PT1H30M" ;

```

```

        :time_coverage_resolution = "PT1S" ;
        :source = "AVHRRMTA_G-NOAA-L1-v2.0, ERA_INTERIM-ECMWF-WSP-
v1.0" ;
        :platform = "Envisat" ;
        :sensor = "AATSR" ;
        :metadata_conventions = "Unidata Dataset Discovery v1.0" ;
        :metadata_link = "http://www.esa-cci.org" ;
        :keywords = "Oceans > Ocean Temperature > Sea Surface
Temperature" ;
        :keywords_vocabulary = "NASA Global Change Master Directory
(GCMD) Science Keywords" ;
        :standard_name_vocabulary = "NetCDF Climate and Forecast
(CF) Metadata Convention" ;
        :geospatial_lat_units = "degrees_north" ;
        :geospatial_lat_resolution = "0.009" ;
        :geospatial_lon_units = "degrees_east" ;
        :geospatial_lon_resolution = "0.009" ;
        :geospatial_vertical_min = -1.e-05f ;
        :geospatial_vertical_max = -1.e-05f ;
        :acknowledgment = "Funded by ESA" ;
        :creator_name = "ESA SST CCI" ;
        :creator_email = "science.leader@esa-sst-cci.org" ;
        :creator_url = "http://www.esa-sst-cci.org" ;
        :creator_processing_institution = "These data were produced
at <institution> as part of the ESA SST CCI project."
        :project = "European Space Agency Sea Surface Temperature
Climate Change Initiative" ;
        :publisher_name = "ESACCI" ;
        :publisher_url = "http://www.esa-sst-cci.org" ;
        :publisher_email = "science.leader@esa-SST-cci.org" ;
        :comment = "See the comment attributes to the
sea_surface_temperature and sea_surface_temperature_depth variables for
information about uncertainty estimates." ;
        :northernmost_latitude = 90.f ;
        :southernmost_latitude = -90.f ;
        :easternmost_longitude = -180.f ;
        :westernmost_longitude = 180.f ;
        :geospatial_lat_max = 90.f ;
        :geospatial_lat_min = -90.f ;
        :geospatial_lon_max = -180.f ;
        :geospatial_lon_min = 180.f ;
        :processing_level = "L2P" ;
        :cdm_data_type = "swath" ;
        :product_specification_version = "SST_CCI-PSD-UKMO-201-
Issue-1-signed" ;
        :full_file = "20100701000000-ESACCI-L2P_GHRSSST-SSTskin-
AVHRRMTA_G-v02.0-fv01.0.nc" ;
        :full_file_uuid = "D7A88FA8-7421-4039-807C-B551D638EDC6" ;
        :adj_file = "20100701000000-ESACCI-L2P_GHRSSST-SSTskin-AATSR-
DM_Adj_<Algorithm>00.2m_10.30-v02.0-fv01.0.nc" ;
        :adj_file_uuid = "F7A88FA8-7421-4039-807C-B551D638EDC6" ;
    }

```

B.2.2.2 L2P auxiliary file containing SST adjustments

```

netcdf 20100701000000-ESACCI-L2P_GHRSSST-SSTskin-
AVHRRMTA_G_Adj_Algorithm_00.2m_10.30-v02.0-fv01.0 {
dimensions:
    ni = 512 ;

```



```

    nj = 28000 ;
    time = 1 ;
variables:
    short sst_adjustments(time, nj, ni) ;
        sst_adjustments:_FillValue = -32768s ;
        sst_adjustments:units = "kelvin" ;
        sst_adjustments:scale_factor = 0.01f ;
        sst_adjustments:add_offset = 0.f ;
        sst_adjustments:long_name = "Adjustment for SSTs to make
them correspond to 10.30 local time and depth of 20 cm" ;
    short adjustment_uncertainty(time, nj, ni) ;
        adjustment_uncertainty:_FillValue = -32768s ;
        adjustment_uncertainty:units = "kelvin" ;
        adjustment_uncertainty:scale_factor = 0.01f ;
        adjustment_uncertainty:add_offset = 0.f ;
        adjustment_uncertainty:long_name = "Time and depth
adjustment uncertainty" ;
        adjustment_uncertainty:correlation_length_scale = "100 km" ;
        adjustment_uncertainty:correlation_time_scale = "1 day" ;

// global attributes:
    :Conventions = "CF-1.5, Unidata Observation Dataset v1.0" ;
    :title = "ESA SST CCI AATSR L2P product" ;
    :summary = "AATSR L2P product from the ESA SST CCI project,
produced using <algorithm name>." ;
    :references = "Insert published or web-based references that
describe the data or methods used to produce them" ;
    :institution = "ESACCI" ;
    :history = "SST CCI processor XXX.YY" ;
    :licence = "TBC" ;
    :id = "AVHRRMTA_G-ESACCI-L2P-v<Version Number>" ;
    :naming_authority = "org.ghrsst" ;
    :product_version = "1.0" ;
    :uuid = "F7A88FA8-7421-4039-807C-B551D638EDC6" ;
    :tracking_id = "F7A88FA8-7421-4039-807C-B551D638EDC6" ;
    :gds_version_id = "2.0" ;
    :netcdf_version_id = "4.1.1" ;
    :date_created = "20110616T152330Z" ;
    :file_quality_level = 3 ;
    :spatial_resolution = "4 km at nadir" ;
    :start_time = "20100701T000000Z" ;
    :time_coverage_start = "20100701T000000Z" ;
    :stop_time = "20100701T013000Z" ;
    :time_coverage_end = "20100701T013000Z" ;
    :time_coverage_duration = "PT1H30M" ;
    :time_coverage_resolution = "PT1S" ;
    :source = "AVHRRMTA_G-L1-v2.0, ERA_INTERIM-ECMWF-WSP-v1.0" ;
    :platform = "Envisat" ;
    :sensor = "AATSR" ;
    :metadata_conventions = "Unidata Dataset Discovery v1.0" ;
    :metadata_link = "http://www.esa-cci.org" ;
    :keywords = "Oceans > Ocean Temperature > Sea Surface
Temperature" ;
    :keywords_vocabulary = "NASA Global Change Master Directory
(GCMD) Science Keywords" ;
    :standard_name_vocabulary = "NetCDF Climate and Forecast
(CF) Metadata Convention" ;
    :geospatial_lat_units = "degrees_north" ;
    :geospatial_lat_resolution = "0.009" ;
    :geospatial_lon_units = "degrees_east" ;

```

```

:geospatial_lon_resolution = "0.009" ;
:geospatial_vertical_min = -0.2f ;
:geospatial_vertical_max = -1.e-05f ;
:acknowledgment = "Funded by ESA" ;
:creator_name = "ESA SST CCI" ;
:creator_email = "science.leader@esa-sst-cci.org" ;
:creator_url = "http://www.esa-sst-cci.org" ;
:creator_processing_institution = "These data were produced
at <institution> as part of the ESA SST CCI project."
:project = "European Space Agency Sea Surface Temperature
Climate Change Initiative" ;
:publisher_name = "ESACCI" ;
:publisher_url = "http://www.esa-sst-cci.org" ;
:publisher_email = "science.leader@esa-SST-cci.org" ;
:comment = "See the comment attributes to the
sea_surface_temperature and sea_surface_temperature_depth variables for
information about uncertainty estimates." ;
:northernmost_latitude = 90.f ;
:southernmost_latitude = -90.f ;
:easternmost_longitude = -180.f ;
:westernmost_longitude = 180.f ;
:geospatial_lat_max = 90.f ;
:geospatial_lat_min = -90.f ;
:geospatial_lon_max = -180.f ;
:geospatial_lon_min = 180.f ;
:processing_level = "L2P" ;
:cdm_data_type = "swath" ;
:product_specification_version = "SST_CCI-PSD-UKMO-201-
Issue-1-signed" ;
:full_file = "20100701000000-ESACCI-L2P_GHRSSST-SSTskin-
AVHRRMTA_G-v02.0-fv01.0.nc" ;
:full_file_uuid = "D7A88FA8-7421-4039-807C-B551D638EDC6" ;
}

```

B.2.2.3 L3U auxiliary file containing SST retrievals

```

netcdf 20100701000000-ESACCI-L3U_GHRSSST-SSTskin-AATSR-
LT_Aux_OE1_Nadir_2_channel-v02.0-fv01.0 {
dimensions:
    lon = 7200 ;
    lat = 3600 ;
    time = UNLIMITED ; // (1 currently)
variables:
    short sea_surface_temperature(time, lat, lon) ;
        sea_surface_temperature:FillValue = -32768s ;
        sea_surface_temperature:units = "kelvin" ;
        sea_surface_temperature:scale_factor = 0.01f ;
        sea_surface_temperature:add_offset = 273.15f ;
        sea_surface_temperature:long_name = "sea surface temperature
retrieved using 2-channels, dual view" ;
        sea_surface_temperature:standard_name =
"sea_surface_skin_temperature" ;
    short l2p_flags(time, lat, lon) ;
        l2p_flags:long_name = "L2P flags" ;
        l2p_flags:flag_meanings = "microwave land ice lake river
spare views channel" ;
        l2p_flags:flag_masks = 1s, 2s, 4s, 8s, 16s, 32s, 64s, 128s ;
    byte quality_level(time, lat, lon) ;
        quality_level:long_name = "quality level of SST pixel" ;

```

```

        quality_level:flag_meanings = "no_data bad_data
worst_quality low_quality acceptable_quality best_quality" ;
        quality_level:flag_values = 0b, 1b, 2b, 3b, 4b, 5b ;
    short large_scale_correlated_uncertainty(time, lat, lon) ;
        large_scale_correlated_uncertainty:_FillValue = -32768s ;
        large_scale_correlated_uncertainty:units = "kelvin" ;
        large_scale_correlated_uncertainty:scale_factor = 0.01f ;
        large_scale_correlated_uncertainty:add_offset = 0.f ;
        large_scale_correlated_uncertainty:long_name = "Uncertainty
from errors likely to be correlated over large scales" ;
    short synoptically_correlated_uncertainty(time, lat, lon) ;
        synoptically_correlated_uncertainty:_FillValue = -32768s ;
        synoptically_correlated_uncertainty:units = "kelvin" ;
        synoptically_correlated_uncertainty:scale_factor = 0.01f ;
        synoptically_correlated_uncertainty:add_offset = 0.f ;
        synoptically_correlated_uncertainty:long_name = "Uncertainty
from errors likely to be correlated over synoptic scales" ;
        synoptically_correlated_uncertainty:correlation_length_scale
= "100 km" ;
        synoptically_correlated_uncertainty:correlation_time_scale =
"1 day" ;
    short uncorrelated_uncertainty(time, lat, lon) ;
        uncorrelated_uncertainty:_FillValue = -32768s ;
        uncorrelated_uncertainty:units = "kelvin" ;
        uncorrelated_uncertainty:scale_factor = 0.01f ;
        uncorrelated_uncertainty:add_offset = 0.f ;
        uncorrelated_uncertainty:long_name = "Uncertainty from
errors likely to be uncorrelated between SSTs" ;

// global attributes:
        :Conventions = "CF-1.5, Unidata Observation Dataset v1.0" ;
        :title = "ESA SST CCI AATSR L3U product" ;
        :summary = "AATSR L3U product from the ESA SST CCI project,
produced using <algorithm name>." ;
        :references = "Insert published or web-based references that
describe the data or methods used to produce them" ;
        :institution = "ESACCI" ;
        :history = "SST CCI processor XXX.YY" ;
        :licence = "TBC" ;
        :id = "AATSR-ESACCI-L3U-v<Version Number>" ;
        :naming_authority = "org.ghrsst" ;
        :product_version = "1.0" ;
        :uuid = "E7A88FA8-7421-4039-807C-B551D638EDC6" ;
        :tracking_id = "E7A88FA8-7421-4039-807C-B551D638EDC6" ;
        :gds_version_id = "2.0" ;
        :netcdf_version_id = "4.1.1" ;
        :date_created = "20110616T152756Z" ;
        :file_quality_level = 3 ;
        :spatial_resolution = "1.1km at nadir" ;
        :start_time = "20100701T000000Z" ;
        :time_coverage_start = "20100701T000000Z" ;
        :stop_time = "20100701T013000Z" ;
        :time_coverage_end = "20100701T013000Z" ;
        :time_coverage_duration = "PT1H30M" ;
        :time_coverage_resolution = "PT1S" ;
        :source = "AATSR-ESA-L1-v2.0, ERA_INTERIM-ECMWF-WSP-v1.0" ;
        :platform = "Envisat" ;
        :sensor = "AATSR" ;
        :metadata_conventions = "Unidata Dataset Discovery v1.0" ;
        :metadata_link = "http://www.esa-cci.org" ;

```

```

        :keywords = "Oceans > Ocean Temperature > Sea Surface
Temperature" ;
        :keywords_vocabulary = "NASA Global Change Master Directory
(GCMD) Science Keywords" ;
        :standard_name_vocabulary = "NetCDF Climate and Forecast
(CF) Metadata Convention" ;
        :geospatial_lat_units = "degrees_north" ;
        :geospatial_lat_resolution = "0.05" ;
        :geospatial_lon_units = "degrees_east" ;
        :geospatial_lon_resolution = "0.05" ;
        :geospatial_vertical_min = -1.e-05f ;
        :geospatial_vertical_max = -1.e-05f ;
        :acknowledgment = "Funded by ESA" ;
        :creator_name = "ESA SST CCI" ;
        :creator_email = "science.leader@esa-sst-cci.org" ;
        :creator_url = "http://www.esa-sst-cci.org" ;
        :creator_processing_institution = "These data were produced
at <institution> as part of the ESA SST CCI project."
        :project = "European Space Agency Sea Surface Temperature
Climate Change Initiative" ;
        :publisher_name = "ESACCI" ;
        :publisher_url = "http://www.esa-sst-cci.org" ;
        :publisher_email = "science.leader@esa-SST-cci.org" ;
        :comment = "See the comment attributes to the
sea_surface_temperature and sea_surface_temperature_depth variables for
information about uncertainty estimates." ;
        :northernmost_latitude = 90.f ;
        :southernmost_latitude = -90.f ;
        :easternmost_longitude = -180.f ;
        :westernmost_longitude = 180.f ;
        :geospatial_lat_max = 90.f ;
        :geospatial_lat_min = -90.f ;
        :geospatial_lon_max = -180.f ;
        :geospatial_lon_min = 180.f ;
        :processing_level = "L3U" ;
        :cdm_data_type = "grid" ;
        :product_specification_version = "SST_CCI-PSD-UKMO-201-
Issue-1-signed" ;
        :full_file = "20100701000000-ESACCI-L3U_GHRSSST-SSTskin-
AATSR-LT-v02.0-fv01.0.nc" ;
        :full_file_uuid = "F7A88FA8-7421-4039-807C-B551D638EDC6" ;
        :adj_file = "20100701000000-ESACCI-L3U_GHRSSST-SSTskin-AATSR-
LT_Adj_<Algorithm>00.2m_10.30-v02.0-fv01.0.nc" ;
    }

```

B.2.2.4 L3U auxiliary file containing SST adjustments

```

netcdf 20100701000000-ESACCI-L3U_GHRSSST-SSTskin-AATSR-
LT_Adj_Algorithm_00.2m_10.30-fv01.0 {
dimensions:
    lon = 7200 ;
    lat = 3600 ;
    time = 1 ;
variables:
    short sst_adjustments(time, lat, lon) ;
    sst_adjustments:_FillValue = -32768s ;
    sst_adjustments:units = "kelvin" ;
    sst_adjustments:scale_factor = 0.01f ;
    sst_adjustments:add_offset = 0.f ;

```

```
        sst_adjustments:long_name = "Adjustment for SSTs to make
them correspond to 10.30 local time and depth of 20 cm" ;
        short adjustment_uncertainty(time, lat, lon) ;
        adjustment_uncertainty:_FillValue = -32768s ;
        adjustment_uncertainty:units = "kelvin" ;
        adjustment_uncertainty:scale_factor = 0.01f ;
        adjustment_uncertainty:add_offset = 0.f ;
        adjustment_uncertainty:long_name = "Time and depth
adjustment uncertainty" ;
        adjustment_uncertainty:correlation_length_scale = "100 km" ;
        adjustment_uncertainty:correlation_time_scale = "1 day" ;

// global attributes:
        :Conventions = "CF-1.5, Unidata Observation Dataset v1.0" ;
        :title = "ESA SST CCI AATSR L3U product" ;
        :summary = "AATSR L3U product from the ESA SST CCI project,
produced using <algorithm name>." ;
        :references = "Insert published or web-based references that
describe the data or methods used to produce them" ;
        :institution = "ESACCI" ;
        :history = "SST CCI processor XXX.YY" ;
        :licence = "TBC" ;
        :id = "AATSR-ESACCI-L3U-v<Version Number>" ;
        :naming_authority = "org.ghrsst" ;
        :product_version = "1.0" ;
        :uuid = "F7A88FA8-7421-4039-807C-B551D638EDC6" ;
        :tracking_id = "F7A88FA8-7421-4039-807C-B551D638EDC6" ;
        :gds_version_id = "2.0" ;
        :netcdf_version_id = "4.1.1" ;
        :date_created = "20110616T152822Z" ;
        :file_quality_level = 3 ;
        :spatial_resolution = "1.1km at nadir" ;
        :start_time = "20100701T000000Z" ;
        :time_coverage_start = "20100701T000000Z" ;
        :stop_time = "20100701T013000Z" ;
        :time_coverage_end = "20100701T013000Z" ;
        :time_coverage_duration = "PT1H30M" ;
        :time_coverage_resolution = "PT1S" ;
        :source = "AATSR-ESA-L1-v2.0, ERA_INTERIM-ECMWF-WSP-v1.0" ;
        :platform = "Envisat" ;
        :sensor = "AATSR" ;
        :metadata_conventions = "Unidata Dataset Discovery v1.0" ;
        :metadata_link = "http://www.esa-cci.org" ;
        :keywords = "Oceans > Ocean Temperature > Sea Surface
Temperature" ;
        :keywords_vocabulary = "NASA Global Change Master Directory
(GCMD) Science Keywords" ;
        :standard_name_vocabulary = "NetCDF Climate and Forecast
(CF) Metadata Convention" ;
        :geospatial_lat_units = "degrees_north" ;
        :geospatial_lat_resolution = "0.05" ;
        :geospatial_lon_units = "degrees_east" ;
        :geospatial_lon_resolution = "0.05" ;
        :geospatial_vertical_min = -0.2f ;
        :geospatial_vertical_max = -1.e-05f ;
        :acknowledgment = "Funded by ESA" ;
        :creator_name = "ESA SST CCI" ;
        :creator_email = "science.leader@esa-sst-cci.org" ;
        :creator_url = "http://www.esa-sst-cci.org" ;
```

```
        :creator_processing_institution = "These data were produced
at <institution> as part of the ESA SST CCI project."
        :project = "European Space Agency Sea Surface Temperature
Climate Change Initiative" ;
        :publisher_name = "ESACCI" ;
        :publisher_url = "http://www.esa-sst-cci.org" ;
        :publisher_email = "science.leader@esa-SST-cci.org" ;
        :comment = "See the comment attributes to the
sea_surface_temperature and sea_surface_temperature_depth variables for
information about uncertainty estimates." ;
        :northernmost_latitude = 90.f ;
        :southernmost_latitude = -90.f ;
        :easternmost_longitude = -180.f ;
        :westernmost_longitude = 180.f ;
        :geospatial_lat_max = 90.f ;
        :geospatial_lat_min = -90.f ;
        :geospatial_lon_max = -180.f ;
        :geospatial_lon_min = 180.f ;
        :processing_level = "L3U" ;
        :cdm_data_type = "grid" ;
        :product_specification_version = "SST_CCI-PSD-UKMO-201-
Issue-1-signed" ;
        :full_file = "20100701000000-ESACCI-L3U_GHRSSST-SSTskin-
AATSR-LT-v02.0-fv01.0.nc" ;
        :full_file_uuid = "D7A88FA8-7421-4039-807C-B551D638EDC6" ;
    }
```