

Perfo radiometre: Mesures 5E vs P2

Study variable	TRO_HUM_RAD_ALG02
Reference variable	TRO_HUM_RAD_ALG06
Missions	AltiKa (<i>al</i>)
Period	[23083.0, 23889.0]

Creation date : 2015/11/18

Contents

A002 - Temporal evolution of differences between both altimetric components	2
A003 - Map of differences between both altimetric components over all the period	3
A004 - Periodogram derived from temporal evolution of altimetric component differences	4
A101 - Temporal evolution of SSH crossovers	6
A102 - Differences between temporal evolution of SSH crossovers	8
A103 - Map of SSH crossovers	9
A104 - Differences between maps of SSH crossovers	10
A105 - Differences between SSH crossovers vs coastal distance	11
A201 - Temporal evolution of Sea Level Anomaly (SLA)	12
A202 - Differences between temporal evolution of Sea Level Anomaly (SLA)	18
A203 - Map of Sea Level Anomaly (SLA) over all the period	20
A204 - Differences between maps of SLA trends	23
A205 - Differences between maps of SLA amplitude and phase	25
A206 - Periodogram derived from temporal evolution of Sea Level Anomaly (SLA)	27
A209 - Differences between maps of SLA variance	30



Diagnostic A002 (mission al)

Name : Temporal evolution of differences between both altimetric components

Input data : Along track altimetric components

Description : The temporal evolution of global statistics (mean, variance, slope) of differences between 2 different standards of a same altimetric component (sea surface height correction, altimeter parameter, orbit) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) . These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Mono-mission analyses





Diagnostic A003 (mission al)

Name : Map of differences between both altimetric components over all the period

Input data : Along track altimetric components

Description : The map of global statistics (mean, standard deviation) of differences between 2 different standards of a same altimetric component (sea surface height correction, altimeter parameter, orbit) are calculated over a given period which is the longer as possible to have obtain reliable statically results. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Mono-mission analyses





Diagnostic A004_a (mission al)

Name : Periodogram derived from temporal evolution of altimetric component differences

Input data : Along track altimetric components

Description : The periodogram derived from temporal and global altimetric component differences is calculated from cycle by cycle monitoring of altimetric component differences (derived from diagnostic A001). It is calculated from the mean or the variance differences. The Periodogram can be calculated for all the periods, but it can be focused on a dedicated period.

Diagnostic type : Mono-mission analyses





Diagnostic A004_b (mission al)

Name : Periodogram derived from temporal evolution of altimetric component differences

Input data : Along track altimetric components

Description : The periodogram derived from temporal and global altimetric component differences is calculated from cycle by cycle monitoring of altimetric component differences (derived from diagnostic A001). It is calculated from the mean or the variance differences. The Periodogram can be calculated for all the periods, but it can be focused on a dedicated period.

Diagnostic type : Mono-mission analyses





Diagnostic A101_a (mission al)

Name : Temporal evolution of SSH crossovers

Input data : Sea Surface Height (SSH) crossovers

Description : The temporal evolution of global statistics (mean, standard deviation) of SSH differences are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Mono-mission analyses





Diagnostic A101_b (mission al)

Name : Temporal evolution of SSH crossovers

Input data : Sea Surface Height (SSH) crossovers

Description : The temporal evolution of global statistics (mean, standard deviation) of SSH differences are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Mono-mission analyses





Diagnostic A102 (mission al)

Name : Differences between temporal evolution of SSH crossovers

Input data : Sea Surface Height (SSH) crossovers

Description : The difference of temporal evolution between the global statistics (mean, standard deviation) of SSH differences are calculated using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Mono-mission analyses





Diagnostic A103 (mission al)

Name : Map of SSH crossovers

Input data : Sea Surface Height (SSH) crossovers

Description : The differences between maps of SSH crossovers differences (mean, variance) are calculated using successively both altimetric components in the SSH calculation. SSH crossovers are the differences between ascending and descending passes for time differences between both passes lower than 10 days (in order to reduce the effect of the oceanic variability).

Diagnostic type : Mono-mission analyses





Diagnostic A105 (mission al)

Name : Differences between SSH crossovers vs coastal distance

Input data : Sea Surface Height (SSH) crossovers

Description : The differences of SSH variances at crossovers are plotted in function of coastal distance, latitudes and longitudes.

Diagnostic type : Mono-mission analyses




Diagnostic A201_a (mission al)

Name : Temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes, or separating North and South hemispheres.

Diagnostic type : Mono-mission analyses





Diagnostic A201_b (mission al)

Name : Temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes, or separating North and South hemispheres.

Diagnostic type : Mono-mission analyses





Diagnostic A201_c (mission al)

Name : Temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes, or separating North and South hemispheres.

Diagnostic type : Mono-mission analyses





Diagnostic A201_d (mission al)

Name : Temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes, or separating North and South hemispheres.

Diagnostic type : Mono-mission analyses




Diagnostic A201_e (mission al)

Name : Temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes, or separating North and South hemispheres.

Diagnostic type : Mono-mission analyses





Diagnostic A201_f (mission al)

Name : Temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The temporal evolution of SLA statistics (mean, standard deviation) are calculated from a cyclic way (altimeter repetivity, daily, weekly, monthly) using successively both altimetric components in the SLA calculation. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements. They are calculated globally, but also separating ascending and descending passes, or separating North and South hemispheres.

Diagnostic type : Mono-mission analyses





Diagnostic A202_a (mission al)

Name : Differences between temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The differences between temporal evolution of SLA are calculated from statistics derived from diagnostic A201 (mean, variance) using 2 different components in the SLA calculation. They are calculated globally, but also separating ascending and descending passes or separating North and South hemispheres.

Diagnostic type : Mono-mission analyses





Diagnostic A202_b (mission al)

Name : Differences between temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The differences between temporal evolution of SLA are calculated from statistics derived from diagnostic A201 (mean, variance) using 2 different components in the SLA calculation. They are calculated globally, but also separating ascending and descending passes or separating North and South hemispheres.

Diagnostic type : Mono-mission analyses





Diagnostic A203_a (mission al)

Name : Map of Sea Level Anomaly (SLA) over all the period

Input data : Along track SLA

Description : The map of global statistics (mean, standard deviation) of SLA are calculated using successively both altimetric components in the SLA calculation over a large period. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Mono-mission analyses



Diagnostic A203_b (mission al)


Name : Map of Sea Level Anomaly (SLA) over all the period

Input data : Along track SLA


Description : The map of global statistics (mean, standard deviation) of SLA are calculated using successively both altimetric components in the SLA calculation over a large period. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Mono-mission analyses

SLA with TRO_HUM_RAD_ALG02 trends : even pass numbers
Mission al, cycles 1 to 23



SLA with TRO_HUM_RAD_ALG06 trends : even pass numbers
Mission al, cycles 1 to 23



Diagnostic A203_c (mission al)


Name : Map of Sea Level Anomaly (SLA) over all the period

Input data : Along track SLA


Description : The map of global statistics (mean, standard deviation) of SLA are calculated using successively both altimetric components in the SLA calculation over a large period. These statistics are calculated from 1 Hz altimetric measurements after removing spurious sea level measurements.

Diagnostic type : Mono-mission analyses

SLA with TRO_HUM_RAD_ALG02 trends : odd pass numbers
Mission al, cycles 1 to 23



SLA with TRO_HUM_RAD_ALG06 trends : odd pass numbers
Mission al, cycles 1 to 23




Diagnostic A204_a (mission al)

Name : Differences between maps of SLA trends

Input data : Along track SLA

Description : The difference of SLA maps (mean, standard deviation, slope) is calculated from maps derived from diagnostic A203 using successively both altimetric components in the SLA calculation over a given period. This can be done globally, or separating in ascending and descending passes (except for SLA Grids).

Diagnostic type : Mono-mission analyses



Diagnostic A204_b (mission al)


Name : Differences between maps of SLA trends

Input data : Along track SLA


Description : The difference of SLA maps (mean, standard deviation, slope) is calculated from maps derived from diagnostic A203 using successively both altimetric components in the SLA calculation over a given period. This can be done globally, or separating in ascending and descending passes (except for SLA Grids).

Diagnostic type : Mono-mission analyses

HUM_RAD_ALG02 trends - SLA with TRO_HUM_RAD_ALG06 trends : even
Mission al, cycles 1 to 23



HUM_RAD_ALG02 trends - SLA with TRO_HUM_RAD_ALG06 trends : odd
Mission al, cycles 1 to 23



Diagnostic A205_a (mission al)


Name : Differences between maps of SLA amplitude and phase

Input data : Along track SLA


Description : The difference of SLA maps (mean, standard deviation, slope) is calculated from maps derived from diagnostic A203 using successively both altimetric components in the SLA calculation over a given period. This can be done globally, or separating in ascending and descending passes (except for SLA Grids).

Diagnostic type : Mono-mission analyses

UM_RAD_ALG02 amplitude - SLA with TRO_HUM_RAD_ALG06 amplitude
Mission al, cycles 1 to 23



O_HUM_RAD_ALG02 phase - SLA with TRO_HUM_RAD_ALG06 phase : a
Mission al, cycles 1 to 23





Diagnostic A205_b (mission al)

Name : Differences between maps of SLA amplitude and phase

Input data : Along track SLA

Description : The difference of SLA maps (mean, standard deviation, slope) is calculated from maps derived from diagnostic A203 using successively both altimetric components in the SLA calculation over a given period. This can be done globally, or separating in ascending and descending passes (except for SLA Grids).

Diagnostic type : Mono-mission analyses





Diagnostic A206_a (mission al)

Name : Periodogram derived from temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The periodogram derived from temporal evolution of SLA (global, northern or southern hemisphere) can be done over all periods or focusing on particular periods, such as annual, semi annual or 60 day signal. Therefore mean of SLA differences are computed (every day or cycle), and time data series are plotted as a periodogram.

Diagnostic type : Mono-mission analyses





Diagnostic A206_b (mission al)

Name : Periodogram derived from temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The periodogram derived from temporal evolution of SLA (global, northern or southern hemisphere) can be done over all periods or focusing on particular periods, such as annual, semi annual or 60 day signal. Therefore mean of SLA differences are computed (every day or cycle), and time data series are plotted as a periodogram.

Diagnostic type : Mono-mission analyses





Diagnostic A206_c (mission al)

Name : Periodogram derived from temporal evolution of Sea Level Anomaly (SLA)

Input data : Along track SLA

Description : The periodogram derived from temporal evolution of SLA (global, northern or southern hemisphere) can be done over all periods or focusing on particular periods, such as annual, semi annual or 60 day signal. Therefore mean of SLA differences are computed (every day or cycle), and time data series are plotted as a periodogram.

Diagnostic type : Mono-mission analyses




Diagnostic A209 (mission al)

Name : Differences between maps of SLA variance

Input data : Along track SLA

Description : The differences between maps of SLA are calculated from the SLA differences (mean, standard deviation) using successively both altimetric components in the SLA calculation.

Diagnostic type : Mono-mission analyses




Diagnostic A210_a (mission al)

Name : Differences between maps of SLA variance for different frequency bands

Input data : Along track SLA

Description : The differences between maps of SLA (variance) are calculated from the mean SLA maps using successively both altimetric components in the SLA calculation filtered to separate high-frequency ($T < 1$ yr), mid-frequency ($1 \text{ yr} < T < 3$ yrs) and low-frequency ($T > 3$ yrs) signals.

Diagnostic type : Mono-mission analyses




Diagnostic A210_b (mission al)

Name : Differences between maps of SLA variance for different frequency bands

Input data : Along track SLA

Description : The differences between maps of SLA (variance) are calculated from the mean SLA maps using successively both altimetric components in the SLA calculation filtered to separate high-frequency ($T < 1$ yr), mid-frequency ($1 \text{ yr} < T < 3$ yrs) and low-frequency ($T > 3$ yrs) signals.

Diagnostic type : Mono-mission analyses



Diagnostic A210_c (mission al)

Name : Differences between maps of SLA variance for different frequency bands

Input data : Along track SLA

Description : The differences between maps of SLA (variance) are calculated from the mean SLA maps using successively both altimetric components in the SLA calculation filtered to separate high-frequency ($T < 1$ yr), mid-frequency ($1 \text{ yr} < T < 3$ yrs) and low-frequency ($T > 3$ yrs) signals.

Diagnostic type : Mono-mission analyses

